



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines


Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>



5

IMPROVED EDITIONS.

SCHOOL ATLASES

By ALEX. KEITH JOHNSTON,

LL.D. F.R.S.E. F.R.G.S. F.G.S.


Geographer to Her Majesty for Scotland, and to the Royal and Physical Atlases, &c.

Map of Scotland 27. A

PHYSICAL GEOGRAPHY

Designs, the elementary facts of this Atlas of Physical Geography, than in the previous works of the form and structure of our Planet, proceed systematically, from a re-

lands, Plains, and Valleys, of the different portions of the Globe, and the distribution of Earthquakes, Volcanoes, Heat, Rain, and Wind—by which its surface is modified—to its actual occupation by the various Races, Families, and Species of Plants, Animals, and Man. A New and Enlarged Edition, containing Twenty Coloured Maps and Letterpress. Imperial Octavo, half-bound, 12s. 6d.



in a Series of Original

ROLOGY, and NATURAL HISTORY. In more simple and elementary manner convey broad and general ideas on the affecting its outer crust. Its plan is to

Lakes, Rivers, Mountain-Chains, Table-

CLASSICAL GEOGRAPHY. A New and Enlarged Edition, 23

Coloured Maps; constructed from the best materials, and embodying the results of the most recent investigations, accompanied by a complete INDEX OF PLACES, in which the proper quantities of the Syllables are marked by T. HARVEY and E. WORSLEY, M.M.A. Oxon. Imperial Octavo, half-bound, 12s. 6d.

OPINIONS OF THE PRESS.

"We have received a new and enlarged edition of the 'School Atlas of Classical Geography,' by Alex. Keith Johnston, with additional maps and names, a complete index, with modern equivalents, and the special attraction of Mr Gladstone's co-operation, who not only placed at the editor's disposal the illustrations to his work on Homer, but enhanced the favour by revising the proof-sheets of the plates and text, as adapted for this Atlas—certainly the most complete work of the kind."—*Spectator*.

"The recently-published edition of Mr A. Keith Johnston's 'Atlas of Classical Geography' is so much enlarged and improved as to be virtually a new work, surpassing anything else of the kind extant, both in utility and beauty. A more complete and in every way satisfactory companion to the study of ancient history and geography need hardly be desired. The maps are most carefully delineated, with a strict accuracy, extending even to the minutest details, the lines being very distinct, and the names, even when necessarily crowded, always easily legible. In short, there is a delicate finish of execution about the whole, which is at once gratifying to the eye and instructive to the mind."—*Athenaeum*.

GENERAL AND DESCRIPTIVE GEOGRAPHY, Exhibiting

the actual and comparative Extent of all the Countries in the World; with their present Political Divisions.

A New and Enlarged Edition, suited to the best Text Books; with Geographical information brought up to the time of publication. Twenty-six Maps, clearly and uniformly printed in colours, with Index. Imperial Octavo, half-bound, 12s. 6d.

ASTRONOMY. A New and Enlarged Edition, 21 Coloured

Plates. With an ELEMENTARY SURVEY OF THE HEAVENS, designed as an accompaniment to this Atlas, by ROBERT GRANT, M.A. LL.D. F.R.S. F.R.A.S., Professor of Astronomy and Director of the Observatory in the University of Glasgow. Imperial Octavo, half-bound, 12s. 6d.

CONTENTS.

PLATE 1. THE CELESTIAL SPHERE—REFRACTION, PARALLAX, ABERRATION—PHASES OF THE MOON—PHASES OF THE INFERIOR PLANETS.—2. DAY AND NIGHT—THE SEASONS—NEAP TIDES—SPRING TIDES.—3. THE SUN AND THE SOLAR SPOTS—THE ZODIACAL LIGHT, &c.—4. MAP OF THE MOON—PHASES OF THE MOON, &c.—5. SOLAR ECLIPSES.—6. LUNAR ECLIPSES.—7 AND 8. THE SOLAR SYSTEM.—9. SATURN, MARS, JUPITER

AND HIS SATELLITES.—10. COMETS.—11. THE STARS, THE MILKY WAY, DOUBLE STARS, COLOURS OF THE STARS.—12. NEBULAE.—13. TRACK OF METEORS 1866—PRISMATIC SPECTRA OF DO.—14. SPECTRUM ANALYSIS APPLIED TO THE SUN AND STARS.—15 TO 20 INCLUSIVE. SIX MAPS OF THE STARS—ECLIPTIC CHARTS—APPARENT PATHS OF THE PLANETS VENUS, MARS, ETC.

"They are as superior to all School Atlases within our knowledge, as were the larger works of the same Author in advance of those that preceded them."—*Educational Times*.

"Decidedly the best School Atlases we have ever seen."—*English Journal of Education*.

"The *Physical Atlas* seems to us particularly well executed. . . . The last generation had no such help to learning as is afforded in these excellent elementary Maps. The *Classical Atlas* is a great improvement on what has usually gone by that name; not only is it fuller, but in some cases it gives the same country more than once in different periods of time. Thus it approaches the special value of a historical atlas. . . . The *General Atlas* is wonderfully full and accurate for its scale. . . . Finally, the *Astronomical Atlas*, in which Mr Hind is responsible for the scientific accuracy of the maps, supplies an admitted educational want. No better companion to an elementary astronomical treatise could be found than this cheap and convenient collection of maps."—*Saturday Review*.

ELEMENTARY ATLAS OF GENERAL AND DESCRIPTIVE

GEOGRAPHY, for the use of Junior Classes; including a MAP OF CANAAN and PALESTINE, and a GENERAL INDEX. Price 5s. half-bound, in Octavo.

TWELVE OUTLINE PROJ

with the Plates of Keith Johnston's General S SOUTH AMERICA, ENGLAND, SCOTLAND, IRELAND Projections, 3d. each.

ending in Scale AFRICA, NORTH AMERICA, ONE. Price 2s. 6d. Single

NEW WORKS ON GEOGRAPHY

REV. ALEX.

A MANUAL OF MODERN GEOGRAPHY, PHYSICAL, POLITICAL, AND POLITICAL ECONOMY.

This volume—the result of many years' experience of Teachers, Advanced Classes, Candidates for the B.A. Degree, &c.

ELEMENTS OF MODERN GEOGRAPHY.

TWENTY-SECOND THOUSAND. 1891.

The *Elements* form a careful and complete summary of the same, the river-systems of the globe, the political boundaries, and the results of the latest census, are introduced into many of the best schools.

OUTLINES OF MODERN GEOGRAPHY.

REVISED TO THE PRESENT TIME.

These *Outlines*—in many respects the wants of beginners. The arrangement of details are avoided, the broad outlines of the subject are given, and the most recent changes in political geography are introduced.

FIRST STEPS IN GEOGRAPHY.

pp. 56. Sewed, 4d. In cloth, 6d.

GEOGRAPHY OF THE YOUNG.

CHILDREN. Sewed, 3d.

OPINIONS.

Annual Address of the President of the Royal Geographical Society.—“I must admire the ability and persevering industry of the author in the freshness and originality. In no respect does the author commence his description of the geographical structure. The work is largely sold, indeed, a most useful school-book in open reference.”

Saturday Review.—“It contains a prodigious array of geographical facts, and will be found as useful for reference.”

English Journal of Education.—“Of all the Manuals on Geography that have come under our notice, we place the one whose title is given above in the first rank. For fulness of information, for knowledge of method in arrangement, for the manner in which the details are handled, we know of no work that can, in these respects, compete with Mr Mackay's 'Manual.'”

A. KEITH JOHNSTON, LL.D. F.R.S.E. F.R.G.S., H.M. Geographer for Scotland, Author of the 'Physical Atlas,' &c. &c.—“There is no work of the kind, (the *Elements*) in this or any other language, known to me, which comes so near my ideal of perfection in a school-book, on the important subject of which it treats. In arrangement, style, selection of matter, clearness, and thorough accuracy of statement, it is without a rival; and knowing, as I do, the vast amount of labour and research you bestowed on its production, I trust it will be so appreciated as to insure, by an extensive sale, a well-merited reward.”

G. BICKERTON, Esq., Edinburgh Institution.—“I have been led to form a very high opinion of Mackay's 'Manual of Geography' and 'Elements of Geography,' partly from a careful examination of them, and partly from my experience of the latter as a text-book in the EDINBURGH INSTITUTION. One of their most valuable features is the elaborate Table of River-Basins and Towns, which is given in addition to the ordinary Province or County list, so that a good idea may be obtained by the pupil of the natural as well as the political relationship of the towns in each country. On all matters connected with Physical Geography, Ethnography, Government, &c., the information is full, accurate, and well digested. They are books that can be strongly recommended to the student of geography.”

RICHARD G. GRAHAM, English Master, College for Daughters of Ministers of the Church of Scotland and of Professors in the Scottish Universities.—“No work with which I am acquainted so amply fulfils the conditions of a perfect text-book on the important subject of which it treats, as Dr Mackay's 'Elements of Modern Geography.' In fulness and accuracy of details, in the scientific grouping of facts, combined with clearness and simplicity of statement, it stands alone, and leaves almost nothing to be desired in the way of improvement. Eminent, by reason of this exceptional variety and thoroughness, to meet all the requirements of higher education, it is never without a living interest, which adapts it to the intelligence of ordinary pupils. It is not the least of its merits that its information is abreast of all the latest developments in geographical science, accurately exhibiting both the recent political and territorial changes in Europe, and the many important results of modern travel and research.”

Spectator.—“The

NEW ATLAS FOR TUTORS AND PUPIL-TEACHERS.

THE HANDY ROYAL ATLAS.

BY

ALEX. KEITH JOHNSTON, LL.D., ETC.

AUTHOR OF THE 'ROYAL ATLAS,' THE 'PHYSICAL ATLAS,' ETC.

45 MAPS CLEARLY PRINTED AND CAREFULLY COLOURED, WITH GENERAL INDEX.

A NEW EDITION, BROUGHT UP TO THE PRESENT TIME,
TO WHICH IS ADDED A NORTH POLAR CHART.

Imperial Quarto, price £2, 12s. 6d., half-bound morocco.

This work has been constructed for the purpose of placing in the hands of the public a useful and thoroughly accurate ATLAS of Maps of Modern Geography, in a convenient form, and at a moderate price. It is based on the '**Royal Atlas**,' by the same Author; and, in so far as the scale permits, it comprises many of the excellences which its prototype is acknowledged to possess. The aim has been to make the book strictly what its name implies, a **Handy Atlas**—a valuable substitute for the 'Royal,' where that is too bulky or too expensive to find a place, a needful auxiliary to the junior branches of families, and a *vade mecum* to the tutor and the pupil-teacher.

OPINIONS OF THE PRESS.

"Is probably the best work of the kind now published."—*Times*.

"Not only are the present territorial adjustments duly registered in all these maps, but the latest discoveries in Central Asia, in Africa, and America have been delineated with laborious fidelity. Indeed the ample illustration of recent discovery, and of the great groups of dependencies on the British Crown, renders Dr Johnston's the best of all Atlases for English use."—*Pall Mall Gazette*.

"He has given us in a portable form geography posted to the last discovery and the last Revolution."—*Saturday Review*.

"This is Mr Keith Johnston's admirable Royal Atlas diminished in bulk and scale so as to be, perhaps, fairly entitled to the name of 'handy,' but still not so much diminished but what it constitutes an accurate and useful general Atlas for ordinary households."—*Spectator*.


"Thus we are inclined to give Mr Alexander Keith Johnston every credit for the production of a handy and at the same time accurate and excellent Atlas, and bringing within reach of every one the advantages without the bulk of his admirable Royal Atlas."—*Examiner*.

"The 'Handy Atlas' is thoroughly deserving of its name. Not only does it contain the latest information, but its size and arrangement render it perfect as a book of reference."—*Standard*.


"It presents to the public, in a convenient form and at a very moderate price, a thoroughly accurate Atlas of Modern Geography."—*Observer*.

WILLIAM BLACKWOOD AND SONS, EDINBURGH AND LONDON.

ILLUSTRATIONS OF



CHARTOGRAPHY



Force of the Wind		Rain	
0	Hardly perceptible	0.00	Sierra Leone
10	Gentle pleasant Wind	0.25	Mahabaleswar (India)
20	Pleasant brisk Gale	0.50	Philadelphia
30	Very brisk	1.07	
40	High Wind	2.56	
50	Very high	3.07	
60	A Storm	4.42	
70	A great Storm	6.07	
80		7.87	
90		12.50	
100		17.75	
110		21.49	
120		26.00	
130		31.25	
140		37.50	
150		45.00	
160		53.75	
170		64.00	
180		75.00	
190		87.50	
200		101.25	
210		116.25	
220		132.50	
230		150.00	
240		168.75	
250		188.75	
260		210.00	
270		232.50	
280		256.25	
290		281.25	
300		307.50	

GRAPHY Meteor - Aurora Borealis - Reflection - Snow

SCHOOL ATLAS
OF
PHYSICAL GEOGRAPHY

ILLUSTRATING IN A SERIES OF ORIGINAL DESIGNS

THE ELEMENTARY FACTS OF
GEOLOGY, HYDROGRAPHY, METEOROLOGY
AND NATURAL HISTORY

BY
ALEX. KEITH JOHNSTON

LL.D. F.R.S.E. F.R.G.S. F.G.S.

GEOGRAPHER IN ORDINARY TO HER MAJESTY FOR SCOTLAND; AUTHOR OF THE 'PHYSICAL
ATLAS,' THE 'ROYAL ATLAS,' ETC. ETC.

NEW AND ENLARGED EDITION

WILLIAM BLACKWOOD AND SONS
EDINBURGH AND LONDON
MDCCCLXXIII



PREFACE TO FIRST EDITION.

GEOGRAPHY has of late years assumed that position in the business of Education to which, from its interest and importance, it is so well entitled; and it is now found that, in order to do justice to its claims, it must be taught in a manner more systematic and orderly than was formerly considered necessary. The most obvious means of facilitating its study was the separation of its component parts into the great divisions of PHYSICAL or NATURAL, and GENERAL and DESCRIPTIVE GEOGRAPHY; and since public attention was first directed to the advantages of this distinction by the Author's folio Physical Atlas, many elementary treatises have been published for the purposes of general instruction. The subject-matter of each of these will find its appropriate illustration in the following Work, which has been prepared expressly for the purpose of conveying broad and comprehensive views of the Form and Structure of the Earth, and the principal phenomena affecting its outer crust.

Beginning with a representation of the Oceans, Lakes, Rivers, Mountain-Chains, Plains, and Valleys of the different portions of the Globe, it proceeds to the distribution of those elements by which its surface is affected—Earthquakes, Volcanoes, Heat, Wind and Rain; and concludes with the actual occupation of its surface by the different races, families, and species of plants, animals, and man.

The object of PHYSICAL GEOGRAPHY, in treating of inorganic matter, is to represent the Earth in its natural state, divested of the accidental or artificial divisions which have been introduced by man's agency. By a patient study and careful comparison of numerous facts, the knowledge requisite for such a representation has, within a recent period, been greatly extended. Comparative Geography has led to analogous views of the structure of the different Continents similar to those which comparative anatomy has established in relation to the lower animals and man. The bones and arteries of the latter have their representatives in the mountain-chains and river-courses of the former. Asia, in its continental mass, presents a picture of majestic unity; Europe, indented and broken up into numerous peninsulas, exhibits an example of the greatest diversity; while the Western Continent is remarkable for its grand simplicity. Again, contrasting climate and position, we find that Africa, presenting the greater part of its surface to the burning rays of a tropical sun, has all its days and nights of nearly equal length—the diversity of the seasons being almost unknown; while, in the frozen regions around the Poles, night extends its absolute empire, "day disappears with its radiant cortege, or if it shines, it is but the longest meteor of a long night."*

Physical Geography teaches that there is an intimate and reciprocal action of Man on the Earth, and of the earth on man, without attention to which it is not possible to understand the National character or physical development of a people. In the East, his wants being easily supplied, and having no necessity to struggle with nature, Man resigns himself to indifference and fatalism; while in the West, in order that he may live, he must conquer the obstacles which nature opposes to his progress; and hence much of the energy, the perseverance, and the intellectual pre-eminence which characterise the races of the West. The whole character of a nation may, as Dr Arnold observes, be influenced by its geology and physical geography. "Who can wonder," he says, "that the rich and well-watered plain of the Po should be filled with flourishing cities, or that it should have been contended for so often by successive invaders?"† It is the abundance of its coal-mines that gives to England its pre-eminence in the manufacturing world. China is chiefly interesting to us for its cultivation of the tea-plant, the Molucca islands for their spices, and Siberia for its furs. It is owing to the nature of the soil and climate that the Southern States of North America are essentially agricultural; while, from having fewer advantages in this respect, and greater geographical facilities from seaports, the Northern States are almost as essentially manufacturing and commercial.

The eminently suggestive character of the Maps will, it is hoped, enable the intelligent teacher to draw many such contrasts and comparisons. Physical Geography is the history of Nature presented in its most attractive form, the exponent of the wonders which the Almighty Creator has scattered so profusely around us. Few subjects of general education are, therefore, so well fitted to expand and elevate the mind, or satisfy the curiosity of youth.

PREFACE TO THE SECOND EDITION.

IN the present edition four plates appear for the first time,‡ the others have been revised and improved,§ and the text is in great part re-written. For valuable aid in the design and execution of the geological illustrations, the Author is indebted to his friend, Archd. Geikie, Esq., F.R.S., Director of the Geological Survey of Scotland.

EDINBURGH, December 1868.

PREFACE TO THE THIRD EDITION.

AT the suggestion of the Rev. Edward Hale, of Eton College, the cotidal lines have been added to the map, which exhibits the ocean currents; and the opportunity was taken to re-engrave this plate from a more perfect drawing. This forms the chief new feature of the present edition of the Atlas; but the whole of the maps have been subjected to a critical revision, and a large part of the accompanying letterpress has been re-written.

EDINBURGH, February 1873.

* RITTER.

† 'Lectures on Modern History.'

‡ Plates II. and III., Illustrations of Physical Geology; XIII. and XIV., River Systems of the British Isles.

§ The Geological Map of the British Isles has been engraved from a new drawing.

LIST OF PLATES.

	PLATE	DESCRIPTION, PAGE
	1.	ILLUSTRATIONS OF CHARTOGRAPHY AND CLIMATOGRAPHY, 1
PHYSICAL GEOLOGY.	2.	ILLUSTRATIONS OF THE GEOLOGICAL ACTION OF RAIN, STREAMS, ICE, AND SNOW, 7
	3.	ILLUSTRATIONS OF THE GEOLOGICAL ACTION OF SEA AND ICE, 13
	4.	GEOLOGICAL MAP OF THE BRITISH ISLES, 23
	5.	DISTRIBUTION OF EARTHQUAKES AND VOLCANOES OVER THE GLOBE, AND RISING AND SINKING OF CONTINENTS, 17
	6.	THE MOUNTAINS, TABLE-LANDS, PLAINS, & VALLEYS OF EUROPE, 29
TOPOGRAPHY.	7.	" " " " ASIA, 31
	8.	" " " " AFRICA, 32
	9.	" " " " N. AMERICA, 35
	10.	" " " " S. AMERICA, 37
HYDROGRAPHY.	11.	CHART OF THE OCEAN CURRENTS AND COTIDAL LINES, 39
	12.	CHART OF THE RIVER SYSTEMS OF THE WORLD, 43
	13 & 14.	RIVER SYSTEMS, RAINFALL, AND CONTOURS OF THE BRITISH ISLES, 46
METEOROLOGY.	15.	THE DISTRIBUTION OF RAIN AND SNOW OVER THE GLOBE, 49
	16.	CLIMATOLOGICAL CHART, SHOWING THE TEMPERATURE OF THE GLOBE, 50
	17.	MAP OF THE CONSTANT AND PERIODICAL WINDS OF THE GLOBE, 51
NATURAL HISTORY.	18.	MAP OF THE DISTRIBUTION OF VEGETABLE LIFE OVER THE GLOBE, 52
	19.	" " ANIMAL LIFE " " 53
	20.	" " THE DIFFERENT RACES OF MAN " " 54

ATLAS

OF

PHYSICAL GEOGRAPHY.

DESCRIPTION OF THE PLATES.

PLATE I.—Chartography and Olinatography.

CHARTOGRAPHY.

Fig. 1 is that of the Mariner's Compass, distinguishing the true and the magnetic north points. The compass consists of a magnetised needle, bearing a circular card, the outer edge of which is divided into 32 points, half and quarter points, and into 360 degrees. The four chief or *cardinal* points are North, South, East, and West; the E. being towards the right when facing the N. The points in the middle space between two cardinal points are named after both, and written N.E., N.W., S.E., and S.W. A point half-way between one of these last and a cardinal point is named by a compound of the nearest cardinal point and the adjacent point: thus the point between N. and N.E. is termed N.N.E.; the point between E. and N.E. is E.N.E., &c. The points *next* the eight principal points have the word *by* between the name of such point and the next cardinal point: thus the point *next* to N. on the E. side is termed N. *by* E.; that next, on the W. side, N. *by* W., &c. The needle points to the *magnetic* north, which only in a few parts of the world agrees with the true or geographical north, and the difference between them is called the *variation* of the *compass*. The variation is termed easterly when the north point of the needle is drawn to the east of the true north, and westerly when drawn to the westward. The variation differs according to the place of observation on the surface of the earth, and the position of certain slowly changing points of attraction called *magnetic poles*.

Fig. 2 represents a River-Course, with arrows pointing the direction in which the water flows. The *right* bank of a river is that on the right hand of a spectator situated with his back to its source, and the *left* bank that on the left hand.

Fig. 3. A Ship at Anchor. An *Anchorage* is a suitable holding-ground unimpeded with rocks, where ships can conveniently cast anchor. *Sounding* is the mode of ascertaining the depth of water or the nature of the sea-bottom: it is performed by means of a weight attached to a sounding-line, marked at every 10 fathoms, with an intermediate knot at every 5 fathoms, and carrying an apparatus by means of which a sample of the sand or mud of the sea bottom is brought up. The line attached to the

instrument called the *log*, by which the rate of a ship's motion is ascertained, is divided by *knots* into lengths, each of which is in proportion to a geographical mile as half a minute is to an hour. The log being thrown overboard, note is carefully taken of the number of knots which run out in half a minute, and it follows that the vessel is passing through the water at the same number of geographical miles an hour. Hence the term *knot* has come to be used as another name for a nautical mile, so that when a ship is said to make 10 knots, she is progressing at the rate of 10 sea miles an hour. Soundings, or ascertained depths, are marked on sea-charts in fathoms. (See fig. 14.)

Fig. 4. A Lake or Inland Sea is a portion of water surrounded by land. *Woodlands*, or *Selvas*, are lands covered with natural timber. *Llanos* are vast plains of South America, alternately covered with rank vegetation, and reduced to a desert state by drought.

Fig. 5 is that of a Firth or Estuary, an inlet of the sea connected with the mouth of a river. The term firth or frith is sometimes applied to an open channel, as the Pentland Firth, in which case it appears to have originated from the Latin word *fretum*; but as extensively used on the east coast of Scotland, the word is undoubtedly derived from the *fiord* (pronounced *fiurth*) of the opposite shores of Scandinavia.

Fig. 6. Gulf, a recess of the ocean or of a sea into the land. It is often used indifferently with Bay; but the latter term is applied either to a large or small extent of water, while the former is used only to designate an extensive recess of the sea, or opening into the land.

Fig. 7. Bay, a large or limited recess of the ocean, of a sea, or large lake. A sheltered bay is a place of refuge for a ship in a storm; a *Harbour* or *Haven* is a small bay or port for shelter. The mouth or mouths of a river is a term used to designate that portion of country near where a river enters into a lake or estuary, or into the ocean. The French word *embouchure* is often so applied.

Fig. 8. Strait, a narrow passage of water between two continents or islands, or the entrance from the ocean to a gulf or lake. It is sometimes called a *channel* or a *sound*, and is often erroneously written "Straits." (See also fig. 6.)

Fig. 9. Delta, from the Greek Δ , is a term applied to alluvial tracts between the bifurcating branches of a river. The best example of a delta is that at the mouth of the Nile, shown in the plate. An *Isthmus* is a narrow neck of land between two seas, joining two peninsulas or other portions of land. The Suez Canal, cut across the former Isthmus, has geographically made Africa a vast island. A *Cataract* is a sudden fall of a large body of water from a considerable height; the fall of a smaller body of water is called a *Cascade*. A *Rapid* is produced by a gradual declivity in the bed of a river. Rapids are sometimes navigable.

Fig. 10. A Promontory is a more or less tapering projection of the land into an ocean, sea, or lake. *Cape* is a part of the coast extending into the water beyond the general shore-line, as in fig. 6; or the point of a promontory, as in fig. 11; it is also termed a *Point*, *Head*, *Ness* or *Naze*, and *Mull*. *Headland* is a high bluff portion of the coast not projecting far into the sea. *Archipelago*, a sea interspersed with many islands, as the *Ægean Sea*. *Breakers*, a name applied to rocks or banks which impede the motion of waves, and cause them to *break* or foam.

Fig. 11. Peninsula, a portion of land almost surrounded by water. When the neck of land which joins a peninsula to other land is narrow, it is called an *Isthmus*.

Fig. 12. Desert, a tract of sterile land, covered with sand or loose stones; the term is often applied to uncultivated wastes not devoid of vegetation. *Oasis* is an isolated spot in a desert, where perennial vegetation is supported by springs or by Artesian wells. *Marshes* occur in all situations—on heights, in valleys, in woods, or on the sea-shore; they are either of fresh or salt water. Some are constantly under water, while others are periodically dry. They often emit noxious exhalations termed malaria, which render the air unhealthy, and produce fevers.

Fig. 13. A Chain, or range, of mountains, is a series of elevations linked together continuously or closely, so that their length greatly exceeds their breadth: the *Crest* is the highest part of the range. *Peak* is the term applied to the conical or pointed summit of a mountain. A *Table-land*, or *Plateau*, is a portion of the surface of the earth elevated above the general level of the region in which it is situated: it may be entirely surrounded by mountains or cliffs, or only bounded on one side by higher land. A *Water-parting* is the ridge-line, from which waters flow in different directions. A water-parting may either be elevated, as a mountain-top or the crest of a range of mountains, or only a slight undulation above the general level. The *Watershed* is the slope down which the tributary streams flow from the water-parting to the main *channel* or *course* of a river. A *River-basin* is a hollow, enclosed at its rim by a water-parting; the sides are formed by the watershed, and the lowest part is occupied by the channel of the river. A *Portage* is that portion of the water-parting over which goods or boats may be transported from one river-basin to another.

Fig. 14. Hills or Mountains are represented on maps by means of shading, which is, or ought to be, darker in proportion as the height and slope are greater. The shading is produced by lines, or *hachures*, placed alongside of each other, or by graduated tints. The scale in the diagram shows the requisite depth of shadow necessary to represent any elevation or declivity from a plain at the level of the sea to a slope of 45° . *Roads*, or *Roadstead*, an open anchorage-ground, where ships are sheltered under lea of the land. *Sandbank* is an accumulation of sand under water, diminishing the depth of the sea, or, it may be, appearing above water at certain stages of the tide. *Surf* is the name applied to waves which break upon a flat shore, or a sandbank (termed *bar* when it stretches across the mouth of a river or estuary), or over a *reef* of rocks or coral. The character of a coast-line, whether bold or flat, is shown on charts by signs indicating rocks, cliffs, dunes, sand, mud, shingle, &c.

Fig. 15. The different forms of mountains, cliffs, &c., are here shown in profile. In the upper corner is a topographical plan of a *Glacier*, showing the descent of frozen snow; and a *Moraine*, or accumulation of stones brought down by the glacier, at the bottom.

Fig. 16 (upper section) shows the method of oblique hill-shading adopted in the new survey of Switzerland, and other Continental maps. The light is here represented as coming from one side only, thereby increasing the effect of relief. In vertical shading (lower section) the light is supposed to fall perpendicularly. According to this method—which is adopted in the trigonometrical surveys of England and France—the effect is less picturesque, but more accurate. In Fig. 16 (lower section) *Volcanic*

peaks are represented as they would be seen vertically, from a great elevation; and below, the same peaks as seen horizontally from the surface of the earth. (See Cone of a Volcano in Action, fig. 21, lower section.)

Fig. 17. *Contouring* is a method of representing, by lines on a flat surface, all those portions of a mountain or any high land which are of equal elevation. The shaded section is supposed to be a rock in the ocean, the bottom of which, *o k*, is on a level with the surface of the sea at low water: as the tide rises, the line of level, or *contour-line*, is successively raised to 1 *g*, 2 *f*, &c. The lines under the section show the projection of the contours on a flat surface.

Fig. 18. Plan of an isolated mountain, seen vertically, with contours shaded according to the scale, fig. 14. The sections show the outlines of its several sides, with their elevations in feet, and the method of shading.

Fig. 19. A portion of Palestine, with the valley of the Dead Sea, to convey an idea of the method of representing absolute height and depth, vertically and in profile.

Fig. 20. Atolls, or lagoon islands, are circular reefs of coral formation rising out of the sea, and enclosing a lagoon. Other forms of coral reefs are shown in fig. 21.

CLIMATOGRAPHY.

The names of the objects represented in the view of atmospheric phenomena are given by references at the bottom of the Plate.

Atmospheric Refraction.—When a stratum of the atmosphere, next the earth, is accidentally expanded or condensed, distant objects, instead of being elevated, are depressed; and occasionally, from two such strata of different densities being placed together, the objects are elevated by one of them and depressed by the other, and they appear double, one of the images being *direct* and the other *inverted*. The lower figure represents a ship with its masts and rigging: it is on the horizon, and is the actual vessel. Immediately above is an *inverted* image of the vessel, quite distinct in all its details. Owing to the great amount of refraction in the Arctic regions, the sun is seen earlier above the horizon, and disappears later than in lower latitudes—thus shortening, by several days, the dreary polar winters.

Clouds are vapours rendered visible by their condensation. They float in the atmosphere at an elevation varying from a quarter of a mile to nine or ten miles above the surface of the earth. Clouds are classified and divided into seven kinds, three of which, called the cirrus, cumulus, and stratus, are simple; and four—the cirro-cumulus, cirro-stratus, cumulo-stratus, and cumulo-cirro-stratus or nimbus—are compound.

1. The *Cirrus* (called by sailors the “cat’s tail”) is formed of thin parallel filaments, sometimes resembling a brush, at other times woolly hair or light network. It is the least dense of all clouds, and attains the greatest elevation and the greatest variety of form. The fine particles of which this cloud is composed are supposed to be minute crystals of ice or snow-flakes.

2. *Cumulus* is the name applied to convex or conical heaps of clouds extending upwards from a horizontal base. The structure of the cumulus is usually very dense;

they are formed in the lower regions of the atmosphere, and are carried along the current nearest the earth. They often resemble bales of cotton, or distant hills covered with snow.

3. The *Stratus* appears in the form of a horizontal band, which generally forms at sunset and disappears at sunrise. It is a continuous sheet of cloud increasing from below upwards; and being the lowest kind of cloud, its lower surface usually rests on the earth.

4. The *Cirro-cumulus* is formed by the cirrus-cloud, by the breaking up and collapsing of its fibres into small rounded masses, separated by intervals of clear sky. It occurs in dry, warm, summer weather, when it is known as the "*mackerel sky*."

5. The *Cirro-stratus* is a combination of the cirrus and stratus, consisting of slightly-inclined masses, dense in the middle and thin towards the edges. In form and position it often resembles shoals of fishes. It is a precursor of storms.

6. The *Cumulo-stratus* is formed by the blending of the cirro-stratus and the cumulus. It appears when the cumulus is surrounded by small fleecy clouds, just before rain begins to fall, or on the approach of a thunderstorm.

7. *Cumulo-cirro-stratus* or *Nimbus* is the *rain-cloud*, or system of clouds from which rain is falling. It first assumes a black or bluish-black colour, but changes to grey when rain begins to fall.

A *Rainbow* is formed by the sun's rays falling on minute drops of water, as those of rain, or the spray from a fountain, which, on the opposite region of the sky, are refracted into prismatic colours, and reflected to the eye of a person placed with his back to the sun. Lunar rainbows are produced in the same manner by the light of the moon, but the bow, owing to the feeble illuminating power of the moon, is usually colourless.

Lightning.—If, by the sudden precipitation of the vapour of water in the atmosphere, a certain quantity of electricity is disengaged, then a spark is emitted, which passes from one cloud to another, or from a cloud to the earth, and the zigzag form of its passage is caused by the unequal conducting power of different portions of the air. The colour of lightning is generally a dazzling white, but it sometimes has a violet tinge. The noise of thunder, which follows lightning, is caused by the displacement of air by the explosion of the electricity, and the irruption of that which succeeds it in filling up the vacuum.

Waterspout.—This remarkable meteorological phenomenon is generally observed at sea, but sometimes also over sheets of fresh water. From a dense cloud a conical pillar descends tapering with the apex downwards. When over the sea, there are apparently two inverted cones—one projecting from the cloud, and the other from the water below it. Waterspouts originate in adjacent strata of air of different temperatures, running in opposite directions in the upper regions of the atmosphere, condensing the vapour and imparting to it a whirling motion. The water of the sea is not sucked up by a waterspout, as is commonly supposed, only the spray is carried up by the whirling vortex. The water poured out from waterspouts is always fresh, or nearly so.

The *Aurora Borealis* is a luminous electrical phenomenon of great beauty, seen in the northern part of the heavens. It generally appears soon after sunset as a luminous arch, which spreads over the celestial hemisphere and speedily assumes a pyramidal

form, with shooting columns of light on all sides, but mostly towards the zenith. Its forms subside and reappear with a flitting motion, and sometimes, after spreading over the heavens, the aurora bursts into a splendid display of variegated colours. The aurora which appears near the south pole is called *Aurora Australis*. The auroral display is occasionally so extensive that it has been seen at the same time in Europe and America, and in the northern hemisphere as near the Equator as the West Indies. The generally accepted theory of the origin of the aurora, is that it is an electrical or magnetic discharge from the poles of the globe, and this view is supported by the observed movements and agitation of the magnetic needle during the appearance; but the theory has been advocated that the aurora is simply the reflection in the heavens of the oblique rays of the sun, from an illuminated portion of the earth's surface adjacent. Where the reflected surface is frozen, the ice crystals may divide the rays, and give rise to the varied hues of the aurora; whilst the constant movement of the mirrored land or water beneath the sun, due to the rotation of the earth, may give the fitful changes of the light.

Meteors.—Igneous meteors, comprising fire balls and shooting-stars, are luminous bodies which suddenly appear in the sky, generally at a great height above the earth, moving through the heavens with immense velocity, in a direction inclined to the horizon. After shining with great splendour for a few seconds, they sometimes explode with a loud noise. The shooting-stars, which appear in vast numbers from the 9th to the 14th November, about the 11th August, and at other stated times, are supposed to form a group of bodies which revolves about the sun in an elliptical orbit, and which, in passing through the aphelion, come in contact with the atmosphere of the earth with a velocity of about 30 miles a second, when they become ignited and are consumed.

Snow is moisture in a frozen state which falls from the clouds when the temperature of the air containing aqueous vapour is at or below the freezing-point, 32° Fahr. If snow, in falling from the upper atmosphere, passes into strata of a higher temperature, it will be melted before it arrives at the earth's surface; hence it never snows in the torrid zone, nor in the temperate zone during the heat of summer, except on the tops and sides of very high mountains.

The *Barometer* is an instrument contrived to measure the height of a column of mercury supported by the pressure of the atmosphere. The scale attached to the glass tube, on the left of the diagram, shows how much the mercury falls, for every 5000 feet, in ascending into the atmosphere: thus, supposing the temperature uniform, and the height of mercury in the tube to be 30 inches at the level of the sea, it would fall to 20½ inches at 10,000 feet, or to 9½ inches, if carried to a height of 30000 feet above that level. In measuring the height of mountains with the barometer, correction for difference of temperature between the base and summit, at the time of the observation, forms an important element in the calculation. The *Thermometer* is divided into degrees, according to the Fahrenheit, Centigrade, and Reaumur scales—the first of which is used in this country, and the two latter on the Continent. The temperature at which water boils at different elevations, depending on the pressure of the atmosphere, is used as a means of ascertaining the heights of mountains. Thus,



as shown in the Plate, water boils at a temperature of 212° at the level of the sea, when the pressure is about 30 inches. If it were to boil at a temperature of 172° , it would indicate an elevation of 26,000 feet above that level.

In the scale on the right side of the Plate the different kinds of winds are named according to their estimated velocity in miles per hour. Thus, 50 miles per hour indicates a storm, and 80 miles a hurricane. The scale also shows the calculated amount of pressure, in pounds' weight, exercised by each on a square foot of surface. Thus a velocity of 50 miles per hour gives a pressure of 12'3 pounds on a square foot of surface. The scale on the extreme right of the Plate gives the ascertained amount of rain which falls in different places, showing the remarkable increase of quantity in proceeding from temperate to tropical countries: for while at London the annual fall is about 20 inches, in some parts of the Western Ghauts Mountains of India it amounts to 300 inches, or fifteen times as much.

PLATE II. A.—Illustrations of the Action of Rain and Streams.

Over the surface of the whole globe there is everywhere in progress a process of disintegration and decay. From the tops of the highest mountains down to the shores of the sea, the rocks which come to the surface, even the hardest and most compact, are undergoing a gradual destruction. This is effected by a number of agents, chief among which are rain, streams, frost, glaciers, icebergs, and the sea. The result of the combined action of these different forces is the constant removal of solid material from off the surface of the land. The rocks decay, the mouldering parts are washed off, new surfaces are exposed to the same destruction, and give place in turn to others. This process of waste is known in geology by the name of *denudation*—that is, the denuding or laying bare of rock by the removal of other portions of rock which once covered it. In Plates II. and III. A are given some illustrations of the chief forms under which denudation is carried on. The figures from 1 to 9a on Plate II. show the action of *rain* and *streams*, the left-hand column (1 to 6) representing the destructive effects, and the right-hand column (7 to 9a) the reproductive effects, of that action.

Destructive Effects.—Every year, in such a country as Britain, there falls a quantity of rain which, if collected together, would be sufficient to cover the whole surface to a depth of from 2 to 3 feet. Part of this rain sinks underground, and, after a subterranean circulation, reappears in the form of springs. Another and much smaller portion finds its way, by brooks, rivulets, and rivers, into the sea. When the moisture falls upon the land as rain it is nearly pure water, but when, after flowing over that surface, it comes down to the sea, it is loaded with mineral substances, either in mechanical suspension, such as fine mud, or in chemical solution, as is the case to a greater or less extent with all spring-water. These mineral ingredients or impurities are derived from the decay of the rocks of which the land is composed. When, therefore, we look at a muddy river entering the sea, we have before us a proof of the great waste which is going on over the surface of the region drained by that river.

In fig. 1 (Plate II.) the first and most direct effect of the fall of rain is seen in the

print of the raindrop upon soft sand or mud. On a sandy or muddy beach the ground, after a shower of rain, is seen to be covered with thousands of little pits, produced by the impact of the rain-drops. If a gale has been blowing at the time, the prints are slanted in the direction towards which the wind blew, and have the sand or mud ridged up on their farther side.

Fig. 2 illustrates the further progress of the same action. A thick mass of earth, with many stones of all sizes, is there shown to occupy the sides and bottom of a valley. The rain has gradually loosened and washed away the earth round each stone, but the latter has acted as a protection to the portion of earth lying immediately underneath it. Hence, as the soil around has been removed, the stones have appeared to rise out of the ground on pillars of earth. But the weather continues to act, though more slowly, upon these pillars. By degrees the support of the stone gives way, and then the column of earth, no longer protected, crumbles away. Instances of this kind on a small scale are occasionally to be seen in Britain, but they occur in great perfection in certain valleys of the Alps.

When the raindrops coalesce they form tiny rills; these unite into runnels, these into brooks, and so on till we come to the broad and stately river. Whether the stream be small or large, it makes a course for itself, which it tends constantly to widen and deepen. This it does by pushing mud, sand, gravel, or boulders along the sides and bottom of the channel. These materials rub down the rocks over which they are rolled, acting as a kind of file, and enabling the streams to cut for themselves a path through even the hardest rocks. In fig. 3 a river is seen to have in this manner excavated a long winding gorge or ravine.

In the carving-out of a river-course, however, much must necessarily depend upon the nature and arrangement of the rocks on which the water has to act. If they are very hard and compact, the rate of erosion may be extremely slow; if changes occur in their arrangement—softer portions, for example, alternating with others offering more resistance—the river may be turned from side to side in frequent curves. Sometimes a trifling obstacle is sufficient to turn a stream aside. A brook flowing along flat ground, for instance, may be deflected by a tree or a stone. Under similar circumstances a river winds in broad serpentine curves, as shown in fig. 5. It tends always to cut through the narrow part of the loops, and would thus shorten and straighten its course were it not at the same time busy making new loops elsewhere.

It often happens that a stream cuts out part of its course by means of waterfalls. This is specially the case with many ravines. A waterfall may be formed in the bed of a stream at the junction of two or more rocks which possess very unequal powers of resistance—such, for instance, as a hard limestone lying upon a soft shale. In fig. 4 a junction of this kind is shown by a section where we see the softer strata exposed to the dash of the spray from the fall, and crumbling therefore under the overlying harder strata, which project and give rise to the perpendicular waterfall. As the shales moulder, the support of the limestone from time to time gives way, and consequently large masses of rock are precipitated into the bed of the stream. But the soft beds underneath are again excavated, and a new outjutting ledge of limestone by degrees appears. In this manner slice after slice is cut off the face of rock behind

the fall; the cataract recedes up the stream, and leaves behind it a deep gorge or ravine. In the figure (4), the end of a lake is seen to the left. When the ravine has cut its way back to the edge of this sheet of water, the waterfall will begin to diminish in height; and in proportion as it diminishes, the level of the lake will be lowered. In the end the waterfall will disappear, the lake, if not too deep, will be drained, and the river, flowing through the bed of the old lake, will escape through the narrow ravine which has been cut by the upward retrogression of the fall. The long deep gorge in which the river Niagara flows has been thus excavated by the backward movement of the famous falls.

That portion of rain which, instead of flowing off by brooks and rivers, sinks into the soil, sometimes produces striking changes upon the surface. Water percolating underground through loose porous strata removes portions of these, and tends to undermine the rocks that lie above. If this should happen on a slope, the overlying mass may be deprived of its support, and slide down or break off, forming what is known as a landslide. One of the most remarkable recorded landslips which have happened in this country is represented in fig. 6. It took place at Axmouth, near Lyme Regis, on the Dorsetshire coast. The cliffs there consist of chalk and green sand, resting upon stiff lias clay, which slopes towards the sea. There had been much rain during the early winter of 1839, and the rocks of the cliffs had become full of water, which, unable to descend beneath the clay at the bottom, flowed seawards along its surface underneath the cliffs. By this means the strata of the cliffs, which had probably been already undermined by the action of springs, lost their support. The ground cracked in many directions, and in particular a deep fissure was formed, three-quarters of a mile long, 100 to 150 feet deep, and 240 feet broad.

Reproductive Effects.—The sediment worn off the land by rains and streams goes to form the material for the formation of new rocks. Some portions of it accumulate in valleys and lakes, but the greater part finds its way into the sea. Fig. 7 indicates the way in which a river sometimes deposits sand and gravel in a valley, and leaves it in the form of terraces on either side. Such river-terraces are of common occurrence. The upper ones are the oldest, for the rivers cut their way down, and leave their *alluvia* to indicate the former level of the watercourses.

Where a stream enters a lake, and the onward motion of its current is checked by the still water, the sediment which it held in suspension is allowed to settle to the bottom. As it gathers year after year, this sediment makes the lake at this part shallower, until a low flat tract of land is perhaps actually gained from the water. In fig. 8 a gain of this kind is indicated at the mouth of a mountain stream. In the figure underneath (fig. 8a), a section is given of the new ground thus formed in the lake. At the bottom lies the hard rock of the district in highly inclined beds. Over this come the sand and gravel, pushed outwards by the stream. Each new addition of sediment, after travelling over the surface of these deposits, is shot gently over their further edge, which slopes away from the shore. Hence the alluvial materials are extended into the lake, very much as a railway embankment is made. They gradually rise towards the surface of the water, and at last, by the additions which they receive in floods, they get above the usual water-line. Vegetation then

takes root, and helps to raise the general surface of the newly-formed ground.

The name given to such accumulations of sediment at the mouths of streams is *Delta*. Originally this term was applied to the great flat alluvial expanse at the mouth of the Nile (fig. 9), from the resemblance of its general outline to that of the Greek letter of the same name. But the word is now widely used to denote all such low alluvial grounds at the mouths of streams, whether large or small, and whether they have the delta shape or not.

The delta of the Nile (fig. 9) is one of the most symmetrical upon the globe. Its base next the Mediterranean is upwards of 200 miles long, and its length from the sea to Cairo 100 miles. Over this wide expanse of flat alluvial land the river sends innumerable branching channels, the two main arms reaching the Mediterranean at Rosetta and Damietta. All this area has been formed by the mud brought down by the river. It is prevented from extending seawards by a strong marine current which washes its base, and sweeps the sediment away into deeper parts of the Mediterranean.

In the case of the Mississippi delta, on the other hand, the sediment is allowed to settle in the Gulf of Mexico, and to push its way seawards in long branching arms. That arrangement is shown in fig. 9a, which represents the outer end of the delta. The mud and silt carried in suspension by that vast river are deposited in the Gulf, on either side of the narrow channels into which the main stream branches. These channels thus flow along narrow alluvial strips of land which are pushed out into the Gulf at the rate of between 200 and 300 feet in a year.

The vast extent of the deltas of the Mississippi, Amazon, Ganges, and other large rivers, shows the enormous amount of *denudation* which has been carried on by the different agencies of atmospheric waste. These alluvial accumulations, hundreds of feet deep and thousands of square miles in extent, have been formed out of the mud, sand, and gravel which have been worn away from the mountains and plains by the action of rains, streams, frosts, &c. Yet they do not by any means represent the whole amount of loss sustained by the land. A large proportion of the sediment is carried into the sea, and there mingles with the marine deposits, to form with them strata which, in after ages, will be elevated into hills and valleys like those from the rocks of which they have been derived.

PLATE II. B.—Illustrations of the Geological Action of Ice and Snow.

When water passes from the fluid into the solid state, as it does in the act of freezing, it assumes a new character as a geological agent, and begins a new series of geological changes not less remarkable than those which it effects when flowing as a liquid over the surface of the earth. Some of the more striking of these changes are illustrated in this Plate.

Frost.—All rocks are more or less traversed with joints, cracks, or other partings, into which water finds its way. When the water contained in these narrow chinks freezes, it expands, and in the act of expansion exerts a great pressure upon the walls

of rock between which it is confined. After repeated freezings and thawings, these walls get gradually pushed farther and farther away from each other. Hence what was at first a mere crack becomes widened by degrees into a gaping rent, and the rock is split open. When this takes place along the face of a cliff, large masses of rock are from time to time detached, and fall to the base of the precipice. Fig. 10 shows a cliff of this kind traversed with perpendicular and transverse joints, of which frost has taken advantage to loosen and detach many large blocks. These are lying piled up at the bottom. There, however, they are exposed anew to the frosts and thaws of winter, and are still further split up. In climates where the temperature in winter sinks below the freezing-point, frost plays an important part in loosening the cohesion of soils and rocks.

Snow.—In mountainous countries where snow falls heavily, it sometimes accumulates in thick masses upon steep slopes. When springs or rain or the melted water from the snow runs down the slope underneath the snow, the hold which the latter has upon the ground is loosened; and its weight lending it impetus, a large mass of snow sometimes slides bodily down the declivity, sweeping trees, rocks, and houses before it, and carrying ruin far down among the vineyards or corn-fields below. This is known as an *avalanche* (fig. 2). The geological changes produced by the direct action of snow are, however, of slight moment when compared with those which are performed when the snow, solidifying into ice, moves down the valleys as massive resistless glaciers.

Glaciers.—In Alpine and Arctic regions there is a varying limit known as the *line of perpetual snow*, or more simply, the *snow-line*. The snow which falls below this line in winter is melted in summer, while the snow which falls on ground higher than this line is only partially dissolved by the sun's heat. If this latter portion, therefore, had no other way of disappearance, it would rapidly accumulate till the mountains and continents were deeply buried under snow. This state of things, however, does not take place. The snow which gathers above the snow-line slides gently into the higher valleys, where, as it moves slowly downward, it becomes pressed into ice, and takes the form of *glaciers*. Where a broad mass of ground rises above the snow-line, the snow presses downward into every available valley, and the glaciers thus produced are the drainage of the snow-fields. A glacier is thus quite comparable to a river. It is indeed a *river of ice*, and carries off the superfluous snow somewhat as a river carries off the superabundant rain. In fig. 13 a sketch is given of part of one of the broad Norwegian snow-fields, and two of the valleys are shown into which the snow-drainage moves, each of them being in consequence occupied by a glacier. Fig. 18 is a map of part of the glacier district of Monte Rosa, and shows how the snow drains down into the valleys, and gradually takes the form of glaciers, which descend far below the limits of the snow-line. The general aspect of a glacier as seen from its lower end is given in fig. 12. On either side rise the mountainous sides of the valley; in the centre, following all the windings of the valley, lies the glacier. In front is the termination of the glacier, where the ice melts away as fast as it is pushed forward, and where from the terminal cave a rushing rapid river takes its rise. Another glacier, which reaches the sea in Spitzbergen, is shown in fig. 14.

The direct geological results of glacier-action are two-fold:—1. The carrying of enormous quantities of debris from higher to lower levels; and, 2. The erosion of the sides and bottom of the valleys down which the ice moves.

1. The transporting power of a glacier is well seen in fig. 12. Along each side of the glacier large quantities of soil, stones, and blocks of rock, falling from the mountain sides upon the ice, are borne onward, and are at last tumbled down in heaps where the ice finally melts. These are called *moraines*, those along the sides of the glacier being *lateral*, and those at its lower end *terminal moraines*. Where two glaciers, issuing from separate valleys, unite into one glacier, the right lateral moraine of the one and the left lateral moraine of the other come together, and form a central ridge or mound of rubbish down the middle of the enlarged glacier. This is known as a *medial moraine*. It is well seen in fig. 12. Upon the surface of the glacier shown in fig. 14 several lines of medial moraine are represented, indicating the union of a number of glaciers. Some moraines are of great size, and prove how great is the amount of waste going on in many mountainous districts.

2. The erosive power of glaciers gives them a high importance as geological agents. The sand, gravel, and stones which get entangled between the bottom of the ice and the rocky floor beneath it act as so many files to scrape, groove, and polish the rocks. Wherever the floor over which a glacier moves can be examined, it is found to present a smoothed and polished surface, which is covered with numerous straight and more or less parallel striæ, varying from the finest hair-lines up to grooves like the ruts of cart-wheels. This kind of surface is not, however, confined to the bottom of the valley, but can be seen along the sides where the ice has retreated a little; and, indeed, often for many hundred feet above the present limits of the ice. The character of the striæ is indicated in fig. 19, which represents an ice-scratched stone from the moraine of an alpine glacier. Several sets of striæ are there indicated, the stone having several times shifted the position in which, jammed between the ice and the rock, it had its surface ground down and striated. The smoothed, polished, and striated surface given by the ice to the rocks on the sides and bottom of a valley is shown upon the rock in the foreground of fig. 15. This is known to geologists as a *roche moutonnée*. Two *perched blocks*, or large stones transported and left by the glacier, are also represented.

The extent to which a glacier is ceaselessly eroding the rocks over which it moves is strikingly proved by the muddy nature of all streams which issue from the lower end of glaciers. The mud is derived from the constant grinding away of the rocks within reach of the ice. The peculiar smoothed and striated surfaces of rock, only to be effected by glacier-action, can often be traced up mountain-sides many hundred feet above the present glaciers, and far down the valleys below the point where the glaciers now melt. Hence the existing glaciers are only the shrunk remnant of what they once were. It is now well ascertained that a large part of Northern Europe has been in recent geological times covered with ice. Throughout the British Islands the ice-markings abound, not only in the valleys, but even on the sides and summits of the minor hills.

Where a glacier issuing from its own valley crosses another valley, it sometimes

succeeds in damming back the drainage of the latter, and a lake is the consequence. Such has been the case in the valley depicted at fig. 15, where the two parallel lines sweeping along the sides of the hills mark two levels at which the water successively stood, when a barrier of ice stretched across the lower end of the valley, and prevented the water from escaping. They are lines of beach, the upper being the older of the two, and pointing to a long period during which the water, ponded back by the ice, stood at that level. A diminution of the glacier then lowered the level of the lake to the second line. Subsequently the glacier withdrew, the lake disappeared, and the drainage resumed its ancient course. These lines are well marked in some parts of the Scottish Highlands, where they are known as *parallel roads*.

Icebergs.—When, as in Arctic and Antarctic regions, a glacier reaches the sea, it breaks off there into large fragments, which float away and are known as icebergs. The seaward end of a Spitzbergen glacier is seen in fig. 14, and a number of fragments of ice are floating in front of it. In fig. 16 the aspect of an iceberg in mid-ocean is shown. The geological results effected by icebergs consist in the transport and deposit over the sea-bottom of such debris as may have fallen upon the surface of the glacier before it reached the sea, and also in tearing up and wearing away the surface of submarine rocks or mud over which they may be driven. Icebergs have sometimes been seen loaded with rock-rubbish, and sailing away into mid-ocean with this freight from Arctic lands.

Coast Ice.—In the far north the cold of winter suffices to freeze the water of the ocean. The first cake of ice formed along shore is borne up with the tide, and, augmented by renewed freezings, becomes at length a shelf of ice stretching for hundreds of miles along the coast, with a breadth of 120 or 130 feet, and a height varying from 25 or 30 to 60 or 70 feet. It is known in Greenland as the *ice foot*. The thaws of summer disengage millions of tons of rubbish, mud, earth, stones, and large blocks of rock, from the cliffs above. These are strewn upon the ice-foot. When the summer storms come, large portions of the ice-foot are detached, and float out to sea, carrying with them their piles of debris. In time they melt away, and their earthy burden is precipitated to the bottom. By this means the floor of the Arctic seas in many places receives annually large quantities of detritus, borne by coast-ice from the Greenland shores.

PLATE III. A.—Illustrations of Sea-Action.

Among the various geological agencies by which the surface of the earth is continually modified, none presents itself under more varied and striking aspects than the sea. Covering, as it does, two-thirds of the entire surface of the globe, and washing thousands of miles of coast-line in every continent, it claims an important place alike in physical geography and in geology. Its waters are always in motion, and in moving they disturb the sand, gravel, or rocks over which they pass. On the calmest summer-day, when the surface of the ocean is scarcely dimpled by a gentle undulation, we may yet watch the wavelets, as they ripple over the beach, catch up the fine sand, and roll it backwards and forwards as they advance and retire. The particles

of sand, in rubbing against each other, are necessarily rounded and lessened. Now from this weak and hardly noticeable action we may trace a gradually increasing exhibition of force, until we behold the billows of a winter storm driving huge blocks of rock shorewards, and bursting in clouds of foam upon the cliffs.

Whether it be gently or with violence, there is a ceaseless tear and wear in progress all round the margin of the sea. At considerable depths beneath the surface of the ocean there is probably little or no action of this kind; but where waves, tides, and currents can act, as they do when they come against a coast-line, the land suffers a gradual loss. In this process of destruction, it is not merely that the sea itself beats against the land, and wears it down, frosts, springs, rain, acting upon the rocks of the shore, corrode and loosen them, and in this way powerfully aid the waves. In many cases, indeed, the waves do little more than remove the detritus which has been produced by these atmospheric agents of degradation. Much also depends upon the set of the tides, the force and direction of the prevalent winds, as well as on the form of the coast-line, and the nature and arrangement of the rocks against which the sea breaks.

Plate III. A. contains some illustrations of different aspects of sea-action. In fig. 1 the rocks dip down beneath the waves like a kind of natural breakwater. In such a case, it is evident that the breakers must burst upon the slope and roll up towards its summit without doing it much damage. Their action is here reduced to a minimum. In the end, however, they succeed, with the help perhaps of frosts and atmospheric disintegration, in loosing some part of the rocky declivity. Other motions follow, until the slope is planed down.

Fig. 2 shows how the waves advance when, in place of a slope, they have obtained a vertical face of rock to undermine. The pieces of stone detached from the cliff by frost or otherwise fall upon the beach, and are there made use of by the waves. In a gale these stones are lifted by the breakers and swung forwards against the base of the cliffs from which they have fallen. Year after year, as this grinding goes on, the stones, originally rough and angular, get rounded and polished, until they sometimes are not unlike cannon-balls. And indeed they are really missiles, while the breakers which drive them on form a kind of sea-artillery. In many places they have battered the base of a precipice until they have hollowed out a cave, into which the waves rush with their freight of gravel and boulders. In other cases they have so undermined the cliff (as shown in fig. 2), that at last the overhanging portion fails to have sufficient support, and topples down upon the beach.

The next two figures illustrate further stages of this action. In fig. 3 a sandstone cliff full of points has been tunnelled, partly by the waves and partly by rain and frosts. The roof of this passage will crumble away while the sides are being demolished, until in the end the arch falls in, and a detached mass of rock is left in advance of the cliff. This has already been the case with the sandstone pillar on the left hand. Formerly it was a part of the main cliff, but by degrees it has been severed, and year by year its bulk is diminishing, under the combined influences of the atmosphere and the waves. In fig. 4 are shown different stages, from the arched passage to the mere sunken reef or skerry. First a hollow or cave is made upon an out-

jutting mass of cliff; the cave is continued till it pierces the headland, and a tunnel is produced; this is widened and heightened, and at length the roof falls in. A detached sea-stack now rises up in front of the cliff; slowly it loses bulk as centuries pass away, until at last it is reduced to but a tangle-covered rock, over which the sea-swell is ever breaking. Such has been the history of the newly-submerged rocks on the right of the figure. Farther to the left rises a pillar of rock now detached from the headland; then comes the arch which is now being worn away, and lastly the main line of cliff, out of which future sea-stacks and islets remain to be carved as the sea advances inland.

When the waves beat upon hard rock, their rate of progress is necessarily very slow. But when they encounter only sand, clay, or other soft materials, they sometimes march onward with melancholy rapidity. Figs. 5 and 6 represent a remarkable example of such a rapid inroad. In the former figure, the old church of Reculver, on the Kentish coast, is represented as it stood in the year 1781, separated from the sea by a considerable space of ground covered with the churchyard and some buildings. Fig. 6 (Lyell) is from a drawing made in 1834. The intervening ground has all been swept away, human bones have been seen protruding from the cliff under the churchyard, and were it not for the artificial breakwater which has been constructed, the church would probably by this time have wholly disappeared. Similar evidence of the advance of the North Sea is abundant along the eastern coast-line of England. In some places the cliffs consist of soft stratified materials (as in fig. 7). In other cases the cliffs are formed of stiff unstratified boulder-clay (fig. 8). Where, as in the latter case, blocks of hard stone or boulders occur abundantly in the substance of the cliff, these accumulate below, and form a kind of temporary breakwater. Sometimes, and more especially where a river enters the sea, a bank of gravel and sand is thrown up by the waves in front of the land, from which it is separated by a channel of still water (fig. 9). In such instances the waves raise up a barrier against their own advance, and protect land which otherwise could not fail to be destroyed. In reality, however, the materials of these natural breakwaters do not withstand the waves. They travel along the coast line even for many miles, driven by the prevalent direction of the breakers; and at last, where the bank ends, are either driven ashore or swept away seawards.

The general result of the destructive action of the sea is to pare off and level down the margin of the land. As it is only the parts of the sea near the surface—those exposed to disturbance by winds—which display this action to any considerable extent, there is a constant tendency towards the production of a plane, the surface of which shall lie beneath the limit to which the influence of wind-waves extends. This surface is known to geologists by the name of a *plane of marine denudation*. We often see it displayed on a small scale along our rocky shores, as in the diagram (fig. 10), where a series of inclined strata dipping inland has been planed down, the worn edges of the beds forming a kind of level surface under the waves. The result has been achieved, however, on a vast scale in some countries, broken and crumpled rocks having been, under the combined forces of denudation, planed down into wide plateaux or table-lands. Traces of such forms of surface are still to be detected in this

country. In Wales, for example, and among the Scottish mountains, geologists have found relics of ancient table-lands, out of which the present system of valleys and lakes has been excavated.

The amount of solid material which is worn off the land every year by the united efforts of rain, frost, springs, glaciers, rivers, and the sea, cannot but be enormous. This material, however, is not lost: carried out beneath the sea, it is there spread over the bottom in layers of sand, silt, mud, and gravel. Year after year these layers increase in depth and solidity. In after ages, when the bed of the sea comes to be elevated into dry land, these deposits will, to a large extent, have become by that time hard rock, and will form then the hills and valleys of the new land. Such has been the origin of most of the hills and mountains now visible upon the surface of the globe. They consist in great part of sandstones, limestones, or rather strata, which once were deposited as sediment over the floor of the ocean, but which, in the course of many revolutions of the earth's surface, have now come to form hard parts of high and rugged land.

The Atlantic is the only one of the oceans whose bed has been at all explored, and our knowledge of it is still limited to that part of it which lies north of the equator. In no direction does this portion of the ocean bed present the form of a smooth basin from shore to shore, but, on the contrary, a series of great valleys and table-lands, with slopes as steep as any to be found on the land of the globe. This will be evident from a glance at the sections across the basin of the Atlantic (fig. 11), which have been chosen as nearly as possible at right angles to one another; but in looking at them it must be borne in mind that the vertical scale of each has been exaggerated to twenty-five times the reality, without which enlargement the undulations would scarcely be visible. The section (*a*) from Trinity Bay, Newfoundland, to Valentia Island, Ireland, on the line of the telegraph cables, crosses one of the shallower regions of the North Atlantic, and shows probably the most generally level part of the whole ocean-bed. The greatest depth in this section is 2400 fathoms—a sinking which corresponds to the elevation of the highest summits of the Alps. The section (*b*) from Cape Race, Newfoundland, to Cape St Roque, Brazil, shows one of the most uneven parts of the ocean-bed. It passes first over the bank of Newfoundland, which stretches south-east from the island for more than 300 miles, at a depth of less than 100 fathoms, plunges suddenly into the deepest part of the ocean, first to 4000, then to 6600 fathoms—perhaps the greatest depth in the North Atlantic—a depth in which the highest mountain of the Himalayas might be set with its peak still 10,000 feet below the level of the sea; rises again by a steep slope to the plateau which fills the centre of the North Atlantic, culminating in the Azores Islands, and sinks once more before reaching Cape St Roque, in the depths which lie in the narrow part of the ocean between the West African and South American coasts. The section (*c*) from Cape St Roque to Cape Palmas, West Africa, crosses a great ridge in mid-ocean, which possibly extends from St Helena to Ascension and St Paul, having these islands for its summits, and thus dividing the ocean-bed here into two distinct basins.

PLATES III. B & V.—Earthquakes, Volcanoes, and Rising and Sinking of Continents.**EARTHQUAKES.**

Earthquakes and Volcanoes are phenomena intimately connected with a heated condition of the interior of the earth, though the nature of this subterranean heat and force is not yet known; and whether it is universal and constant, or partial and temporary, has not been decided. The high temperatures observed in deep mines and artesian wells in many parts of the globe, point indeed to a general accession of temperature inwards, which, if it progresses uniformly, would, at a depth of between 20 and 60 miles, reach a point at which all known substances are in a state of fusion.

The concussions of earthquakes are communicated from below, and consist in commotions of parts of the earth's surface, sometimes so slight as to be almost imperceptible; at other times producing violent convulsions, which leave lasting memorials of their destructive effects. Earthquakes generally advance in a linear direction, undulating, with a velocity of from 25 to 35 miles in a minute; but sometimes in circles or great ellipses, in which, as from a centre, the vibrations extend with decreasing force towards the circumference. Their duration is often extremely short, the most destructive shocks being over in a moment. The great earthquake of Lisbon in 1755 lasted only six minutes.

Earthquakes are generally accompanied by detonations, the noise of which has been compared to the roll of thunder, the rattling of heavy waggons, the irregular discharge of cannon, or the hollow sound of an exploding mine: sometimes these sounds are heard for a considerable length of time without any shock following; and, on the other hand, cases occur in which the severest shocks are not accompanied by any noise whatever. When the earthquake wave, or that undulating motion of the earth's crust producing the phenomena of the earthquake, extends under the bed of the sea, the latter is always agitated. During the earthquake at Lisbon the sea rose 50 feet above its ordinary level. In the open ocean the effect of an earthquake upon a ship resembles that produced by striking on a rock under water.

An illustration of the effects of an earthquake upon the surface of the ground is given in Plate III. B, fig. 5. Towards the left a rounded building has been shattered, and one portion has sunk down to a much lower level than the remaining part. On the right the ground is represented as having opened in a long ragged fissure. Into such rents houses, trees, human beings, and different animals are sometimes precipitated; and if the rent close again immediately, all trace of the objects engulfed is lost. It may happen, however, that a second shock coming close after the first may reopen the fissure, and even eject the buried objects to the surface again.

DISTRIBUTION OF EARTHQUAKES.—In so far as yet known, these may occur at any part of the earth's surface, excepting, perhaps, in those areas in which a gradual movement of depression is taking place. The portions of the map which are tinted over show the regions which are known to be subject to earthquakes, those parts in which the number and intensity of the shocks are greater being more deeply shaded;

it is probable, however, that the areas of the map which are not tinted at all are left blank, because no record has been kept of earthquakes which may have occurred in them.* No country, whatever the nature of its geological formation, is entirely exempt from such visitations; they have occurred even in the loose alluvial soil of Holland. They appear in all zones and at all seasons; and it is probable that the surface of the globe is continually agitated by concussions in some of its points.

The earthquake district of the *Mediterranean* and of *Central Asia* extends from the Azores and Canaries on the west to Lake Baikal on the east, and probably to the Great Desert of Africa, the Nile delta, and the Desert of Arabia on the south. It is the longest and most regular zone of volcanic action on the globe.

From the earliest times volcanic appearances have been observed on a line extending between the Caspian Sea and the Azores, which is the centre whence the earthquake proceeds, and according to distance from which the shocks decrease in frequency and violence; and it appears that the boundaries of the principal concussions are limited by the more or less parallel position of the Pyrenees, the Alps, Carpathian, and Caucasian mountain-chains, in reference to this central line.

Within this district the *Earthquake of Lisbon* occurred on the 1st November 1755. Suddenly a sound of subterranean thunder was heard, which was immediately succeeded by a violent shock, demolishing the greater part of the city; and within six minutes 60,000 persons perished. Its movement was undulatory, and it is estimated that it travelled at the rate of 20 miles in a minute. The axis of the shock formed a line extending from the coast of Morocco, along the western shore of Portugal, to Cork in Ireland. From Lisbon as a centre the line of devastation extended north to Oporto, and south to Ayamonte. Within this space, the sea, fearfully agitated by the concussion of its bed, caused great destruction. At Cadiz, a wave 60 feet high rolled over the land; at Lisbon, the sea rose 50 feet; at Funchal, in Madeira, it rose 15 feet above the highest water-mark; and at Kinsale, in Ireland, it overflowed the market-place. The space within which this earthquake was observed on land extended in an elliptical form from the Island of Madeira to Abo in Finland, and from the north of Scotland to Sardinia. In Scotland, the water in Lochs Lomond, Katrine, Long, and Ness rose and fell repeatedly to an extent of 2 or 3 feet; and a similar movement of the surface occurred in the lakes of Germany, Switzerland, France, and Scandinavia. The agitation of the waves of the sea extended across the Atlantic to the Antilles, where the islands of Antigua, Barbadoes, Guadeloupe, and Martinique were overflowed; and at the same time an unusual movement was observed on the surface of Lake Ontario.

Italy and its islands have been more frequently visited by earthquakes than any other part of Europe. Pompeii and Herculaneum were thus destroyed in A.D. 63, and sixteen years afterwards, the cities, which had in the meantime been rebuilt, were buried under the ashes and lava ejected from Mount Vesuvius. Since that time

* The student is referred to the "British Association Catalogue of Earthquakes" of the Messrs Mallet, which contains a record of earthquakes which have happened in all parts of the globe from B.C. 1606 to A.D. 1850, and a full discussion of the theories of their cause.

earthquakes have been of frequent occurrence in Lower Italy and Sicily, especially in the eighteenth and nineteenth centuries. The great earthquake of *Calabria*—5th February 1783—is remarkable for the concentration of its force, since, within an area of 500 miles, 200 towns and villages were destroyed, nearly 100,000 of the inhabitants perished, and the face of the country was completely changed. The convulsions extended along the granite range of the Apennines from north to south, but were not perceived to the eastward of that chain.

North of the foregoing district, in Basilicata and Principato Citra, with the Val di Diano as a centre, a series of earthquakes commenced on the 16th December 1857, and continued, with intervals, during January 1858. These exhibited many of the most appalling features of the phenomena; hills were levelled and deep fissures formed. Upwards of a hundred hamlets, villages, and towns were injured, many of them ruined, and nearly 10,000 persons perished.

In 1838 a region of Southern Europe, far to the eastward, was visited by a destructive earthquake, the shock of which was felt at Vienna and Constantinople, in Southern Russia, and the countries of the Lower Danube. Greece and Turkey have also been visited by frequent and very violent earthquakes. Hence this district extends over Asia Minor, Syria, and Palestine, to the Caspian Sea and the Caucasian Mountains, where the country in the vicinity of Teflis has suffered severely, both in earlier and later times. Persia and the countries of the Caspian form the connecting link between this and the *district of Central Asia*, where the course of linear concussion follows the direction of the great mountain-system of Thien Shan or Syan Shan, north of which the eastern portion of the Altai Mountains, the Plutonic fissure of Lake Baikal, and the warm springs of the Orkhon River, form the central line of that volcanic convulsion which destroyed Karakorum at the end of the thirteenth century. The countries of Asia have suffered as much from earthquakes as those of Europe. The great earthquake of Syria, in May, A.D. 526, most severely felt at Antioch, is said to have caused the sacrifice of 250,000 lives; and in the same region, in 1837, a shock occurred over an extent of 500 miles in length by 90 in breadth, by which 6000 persons perished. The earthquake of Cutch, in 1819, caused such alterations that the bed of the eastern branch of the Indus, previously only 1 foot deep, was suddenly depressed to a depth of 20 feet. In 1832, a shock passed through the chain of the Hindoo-Koosh, and extended to Bokhara and Kokand.

Earthquake District of India.—Between the years 1800 and 1842, one hundred and sixty-two earthquakes are recorded. In 1843 twenty-three shocks were experienced, and since then four or five have been observed in India each year. Most of these occurred within the region of the Himalayas, including Cabool, Jellalabad, Cashmere, Nepaul, and Assam. They very seldom happen in the peninsula south of latitude 15°. Some of these were very severe. In 1833 a shock occurred in the plains between Delhi and Chittagong, the undulatory movement of which was from north-west to south-east: in the valley of Nepaul 4000 houses were overturned, 414 persons were killed, and 172 wounded. In November 1842, Calcutta was nearly the centre of an emanation, the shock of which extended to Darjeeling on the north, Chittagong on the east, and Monghyr on the west. From Mount Aboo as a centre, a shock

extended in an easterly and westerly direction, from Surat in the south to Simla in the north, over an area of about 100,000 square miles.

The *Earthquake District of Iceland* probably includes the whole of Great Britain, the north of France, Denmark, and Scandinavia. Its connection with the Mediterranean district is shown by the coincidence as to time in the great eruption of Skaptar Jokul of 1783 with the earthquake of Calabria in the same year. The first-recorded earthquake in England occurred in A.D. 974; since that time slight shocks have been felt in all parts of the British Islands, and latterly they have been very frequent at Comrie, in Perthshire.

In *Africa*, earthquakes appear to be unknown, except in a part of Barbary in the north, the region of the Red Sea in the east, and the Cape Colony in the south. In the latter region only three shocks are recorded in the present century, all of them slight—one in 1809, another in 1835, and the third in August 1846, which latter extended over an area of about 400 square miles.

Earthquakes of America.—Both as regards frequency and extent, these equal the earthquakes of the Old World. They occur chiefly along the coast on the western side of the Andes, and on the northern declivity of the mountains of Venezuela, very few shocks having been recorded as taking place to the eastward of the Andes. Among the most remarkable was the *earthquake of Chile*, 20th February 1835, which was felt everywhere between the Island of Chiloe in the south and Copiapo in the north. All the towns and villages between the parallels of 35° and 38° south were destroyed, and the line of coast was permanently raised. In the great *earthquake of Caracas* in 1812, the concussion, like that of Lisbon, formed an ellipse. It was felt from near Bogota in the west to the Gulf of Paria in the east, and formed part of the district of nearly synchronous earthquakes which occurred between 1811 and 1813. One of the great phenomena of tropical America was the eruption of *Cosiguina*, and the synchronous earthquake of New Granada in 1835. A few weeks after the eruption of this volcano, nearly the whole of New Granada was violently convulsed. The *earthquake of Guadeloupe*, 8th February 1843, is one of the most interesting on record. It was felt over a long narrow zone, from New York to the mouth of the Amazon, 3000 miles; but its most violent concussion was confined to Guadeloupe and Antigua. The town of Point-à-Pitre in the former island was entirely overthrown, and 6000 of its people destroyed. The earth opened in several places, sheets of water spouting from the creeks 100 feet in height; and when the chasms closed, men and other objects were engulfed. The *eastern side of the United States* is a region where, though earthquakes have been frequent, they are seldom violent. More than 100 are estimated to have occurred within the past 200 years in the earlier-peopled districts. The oscillations of the ground in all the Appalachian region have a prevailing north-east and south-west direction. One of the most conspicuous earthquakes in North America was that which occurred in the *Mississippi Valley* in 1812. It consisted of a series of concussions, beginning late in 1811, and lasting at intervals for more than a year. It resulted in the submergence of a vast tract bordering the river on the west. This tract was permanently altered in level; and some of the chasms opened in the earth during the shocks are still visible. On 4th January 1843, an earthquake was

perceptibly felt from Ohio and Iowa to Louisiana and Georgia, and from the western frontiers of Missouri and Arkansas to the Atlantic coast. Its duration was estimated at about two minutes, and its rate of motion 33 miles per minute.

VOLCANOES OR BURNING MOUNTAINS.

These mark certain portions of the earth's surface, where there is a direct communication between the atmosphere outside and highly-heated portions of the interior of the globe. At such points of communication, *lava, ashes, scoriæ, volcanic dust, steam, and gases* are emitted. The gradual accumulation of the solid materials round the orifice gives rise to a volcanic hill. A volcano is consequently more or less conical in form. Its summit is usually occupied by a cup-shaped cavity called a *crater* (at the bottom of which lies the *vent* or *focus of eruption*), and its sides are covered with lava, loose blocks of ejected rock, or deposits of cinders and ashes. Volcanoes are *active, dormant, or extinct*. In the active state their craters give out large volumes of steam, with showers of ashes and red-hot stones, while streams of melted lava, escaping either from the crater or from fissures in the sides of the mountain, pour down the slopes, destroying all trees, vineyards, or houses that may chance to lie in their path. A volcano is said to be *dormant* when no eruptions take place from it, but when the escape of steam and gaseous or sulphurous exhalations shows that the subterranean heat can still make itself manifest at the surface. Vesuvius, for instance, had been dormant from time immemorial, until in the year 63 it suddenly became active, and destroyed the two Roman cities of Herculaneum and Pompeii. An *extinct* volcano is one where the eruption of lava, ashes, and steam, with the other signs of volcanic activity, has long ceased. In Central France and in Germany there are numerous extinct volcanoes, in which the craters remain perfect, though now grass-grown and silent. In the British Islands are found abundant traces of still more ancient volcanoes, the craters of which have long since been worn away; but the lava and ashes ejected by them remain, and form portions of hills, and sometimes even entire hills.

In Plate III. B are given three illustrations of volcanoes. Fig. 1 is a map or plan of Vesuvius, showing the small modern crater in the centre, with the large partially-effaced ancient crater-wall of Somma half encircling it, and the courses of the lava-streams down the sides of the mountain. Fig. 2 gives a view of Vesuvius from the sea. The modern crater here again occupies the centre, emitting a column of smoke, while the older crater of Somma is seen behind the present active cone. Fig. 3 is a drawing of three extinct volcanoes of Central France. In each of them the hill consists of a cone of ashes, with a central crater, one side of which has been broken down by the current of black lava. In each case the lava has risen within the crater until the pressure burst the weakest part of the crater-wall, and the lava flowed out over the plain.

In some regions large quantities of steam and gas are evolved from small orifices, distinct from ordinary volcano craters. Near Naples, a well-known instance of this kind is called the Solfatara—a name which is now very generally applied to the orifices from which discharges of this kind take place. In some cases the emission of steam and water is effected by sudden efforts, and with great force. Of this nature are the Geysers of Iceland. These are hot springs which for a few minutes throw up a column of water and steam with great violence to a height of 100 feet or more, and

then sink into rest for a little till they are ready for a renewed outburst. Fig. 4 shows the eruption of one of these Geysers. Such is the force with which the hot water is driven out that it ascends vertically, even in spite of a strong wind which drives the steam to one side, and allows it to condense and fall in a shower of slanting rain. Stones thrown into the hole from which the water ascends are projected to great heights, and sometimes even broken to pieces by the violence of the discharge.

RISING AND SINKING OF CONTINENTS.

Movements of Elevation.—Gradual, and in some cases imperceptible, changes of the relative levels of land and sea are constantly occurring in countries situated at a distance from active volcanoes, and where violent earthquakes are not known. Within the present century the shores of the Baltic, in Scandinavia, have been gradually elevated from 10 to 14 inches above the former level, and are still apparently on the uprise. By the earthquake of Cutch already referred to, the Ulla Bund, a tract 70 miles in length and 16 in breadth, was upheaved 10 feet, and the Runn of Cutch was sunk so as to create a salt marsh, with an area of nearly 200 miles; and by the earthquake of Chile in 1835, a tract not less than 100,000 square miles was permanently elevated about 6 feet above its former level, and parts of the sea-bottom remained dry at high-water, with beds of shellfish adhering to the rocks on which they grew. There are also evidences of upheaval on the coasts of Denmark, and on the Prussian shores of the Baltic; and recent observation points to regions of elevation on the north-east shores of Siberia, and at the northern extremity of Baffin Bay. Proofs of the rising of the land are furnished by lines of what are known as “raised beaches.” These are terraces of sand and gravel, sometimes containing marine shells, which mark old sea-margins. They run along the coast in many parts of the British Islands, at heights of 25, 40, 70 feet, and more, above the present sea-level, each raised beach or terrace indicating a line at which the land on it stood before it reached its present elevation. In Plate III. B, figs. 8 and 9, two illustrations of these features are given. Fig. 8 is a sketch of a remarkable series of ancient beach-lines near the town of St Andrews (*Chambers*). Fig. 9 shows, in a sectional form, how such lines of beach remain on a sloping surface of land. The highest terrace (*a*) is of course the oldest, and marks a time when the sea stood at that level, or the land was so much lower than it is now. The next in descending order is marked *b*, and shows another line at which the sea stood for a long while after the land had risen from *b* to *a*. As a rule, the lower terraces are best preserved, inasmuch as they are younger, and have been for a shorter time exposed to the tear and wear of the elements.

Movements of Depression.—A slow submergence of the land is seen on the south-west coast of Greenland, where ancient buildings are now covered by the sea, and where the Moravian settlers have been obliged more than once to move inland the poles to which they attach their boats, the old poles remaining beneath the water. The most remarkable example, however, of the slow sinking of a wide area of the earth's crust is furnished by the coral reefs of the Pacific and Indian Oceans. The general character of a coral islet is shown in fig. 6, Plate III. B. As there indicated, it consists of a low ring of land, with a sheet of calm water inside, and the wild ocean

outside. At one time it was supposed that these remarkable circular islands marked the sites of submerged volcanoes. It is now known, however, that they point to a slow sinking of the bed of the ocean. The nature of the evidence which they furnish is illustrated by fig. 7, Plate III. B, which shows three sections, each of which marks a stage in the growth of a coral island. In the first section (A) we see a mass of high land rising out of the sea as a mountainous island. Along its margin reefs of coral are formed by the coral polypes, and are known as *fringing reefs*. A slow subsidence of this part of the earth's crust is, however, in progress. The island sinks gradually, but not too fast to allow the coral reef to be kept up by constant additions to its original height. In this way the distance from the reef to the shore of the island inside increases, and the reef becomes what is known as a *barrier reef* (B). The downward movement still going on, and the coral polypes continuing to keep their building abreast of the waves, the old island at last disappears. On its site there now lies an expanse of calm, pale green water, protected by the circle of reef from the open ocean outside. This central water is the *lagoon*. The sea breaking on the reef detaches fragments of it, and scatters them over the reef, along with the sand and mud produced by the grinding of the fragments. In this way small portions of dry ground are formed, which gradually increase until they more or less encircle the lagoon. Seeds drifted over the ocean come ashore and take root, and thus arise the flat, palm-covered coral islets of the Pacific. The regions in which these coral islands occur point, therefore, to wide-spread movements of depression of the crust of the earth. They are built, in short, upon the tops of a submerged continent.

The map (Plate V.) is intended to give a general view of the distribution of earthquakes and volcanoes over the globe, and of the areas which are now, or which have in comparatively recent geological times been, rising or sinking. It will be seen that, on the one hand, earthquakes and movements of upheaval are for the most part found in or not far removed from volcanic regions; while, on the other, the wider areas of submergence are distant from these regions.

PLATE IV.—Geological Map of the British Isles.

Perhaps no part of the earth's surface comprises within the same extent of area so complete an epitome of the structure of the rocky crust of our planet as is to be found within the compass of the British Islands. The geological formations which in other countries spread out over hundreds and thousands of square leagues, are here brought within such narrow limits that a single English county contains within itself a greater variety of rocks than can be seen in some whole kingdoms abroad. In this way, without quitting his country, a native of Great Britain may often obtain, from an examination of the rocks around him, a more correct and vivid conception of the nature of geological reasoning, and of the grandeur of geological history, than if he travelled for hundreds of miles in some other countries.

In the map, to which the following pages are appended as a short explanation, it will be seen that the colours are arranged without much reference to the ordinary geographical subdivisions, and that they cross counties and mountains and rivers, apparently in defiance of all order and symmetry. Each of these colours denotes a

distinct geological formation or group of rocks. In order to understand the reason why they are placed in such irregular grouping, it may be well to attend to a few preliminary explanations.

In examining what is known as the *crust of the earth*—that is, the portion of the earth which is accessible to our observation, whether at the surface, or in caves, quarries, pits, and mines—geologists have found that there are two great classes of rocks, one called AQUEOUS or SEDIMENTARY, the other IGNEOUS. Between these two divisions there is an intermediate group, known as the METAMORPHIC, of which the affinities are sometimes closest with the aqueous series, and sometimes with the igneous. Now it is of these rocks that the crust of the earth is formed, and consequently it is these same rocks which we have to study when we wish to become acquainted with the geological history of the British Isles.

The AQUEOUS rocks, as their name denotes, have resulted from the operation of water. In short, they are simply sediment which has been collected under water and deposited there, layer after layer, just as gravel, sand, and mud are accumulated under water in every pool, river, lake, and sea at the present day. The circumstance that this deposition has taken place in layers, seams, beds, or strata, according as the motion of the currents or the supply of sediment varied, has given rise to the term *stratified*, as applied to these successive sheets of different kinds of sediment piled over each other. As the ocean covers by far the larger part of the globe, so the sedimentary accumulations now forming must be chiefly marine, seeing that not only do the tides and currents carry out sediment from the shore, but streams and rivers are everywhere bearing enormous quantities of sand and mud from the land to the sea. So also in past time the greater part of the debris worn away by rains, springs, frosts, rivers, and breakers, from the surface of the land, must have been laid down under the sea. Hence by far the larger number of the subdivisions into which the aqueous rocks have been grouped are of *marine* origin. In further examining the crust of our planet, we find that these aqueous rocks, which were once ordinary gravel, sand, or mud, contain the remains of both plants and animals, just as we see similar remains carried down by rivers or swept out by tides, and deposited among ordinary sedimentary accumulations. But as our investigation advances, we begin to perceive that, in the great majority of cases, neither the plants nor the animals are quite the same as those which live at the present time. And the older the group of aqueous rocks, the further removed are their *fossils* or *organic remains* from the forms of the living world of to-day. Tracing out these differences, we gradually learn that in the rocky crust on which we live and move there is a history of life and change; that from the oldest strata preserved to us up through all the others which have been successively piled over them, there can be traced an onward and upward progression, race after race of plants and animals succeeding each other, approaching nearer and closer to the existing types, until at last we reach the present time, where, as the crowning act of the long history, Man takes the lead in creation.

Arranged in order of time,—that is, in the order in which they have successively been deposited during the past history of the globe,—the aqueous rocks of the British Islands are—

POST-TERTIARY PERIOD.

RECENT.	{	Modern alluvium, peat, and raised beaches.
POST-PLIOCENE.	{	Ancient alluvia of the Thames, Ouse, etc. Cave-deposits. Glacial drift.

TERTIARY OR CAINOZOIC PERIOD.

PLIOCENE.	Older. Newer.	{	Norwich crag.
		{	Red crag.
		{	Coraline crag.
MIOCENE.		{	Leaf-beds of Mull, basalt of Antrim, lignite of Bovey Tracy.
EOCENE.	Lower. Middle. Upper.	{	Upper part of fluvi-marine beds of Isle of Wight.
		{	Lower part of fluvi-marine beds of Isle of Wight.
		{	Bagshot beds.
		{	London clay.
		{	Woolwich and Reading beds.
		{	Thanet sands.

SECONDARY OR MESOZOIC PERIOD.

CRETACEOUS.	Upper.	{	Chalk.
		{	Upper Greensand and Gault.
	Lower.	{	Lower Greensand.
WEALDEN and PURBECK.		{	Weald clay and Hastings sand.
		{	Purbeck beds.
OOLITIC or JURASSIC.	Upper.	{	Portland beds.
	Middle.	{	Kimmeridge clay.
	Oolite.	{	Coral rag, or Coralline oolite.
		{	Oxford clay.
		{	Cornbrash and Forest Marble Group.
	Lower.	{	Great or Bath oolite.
	Oolite.	{	Fullers' earth.
		{	Inferior oolite.
	Lias.	{	Sands, Upper Lias clay.
		{	Marlstone.
		{	Lower Lias clay.

NEW RED SANDSTONE or TRIASSIC.	Keuper.	{ Penarth or Rhætic beds. Red marls, with rock-salt and gypsum. White and brown sandstones.
	Bunter.	{ Upper red and mottled sandstones. Pebble beds. Lower red and mottled sandstones.
PRIMARY OR PALÆOZOIC PERIOD.		
PERMIAN.		{ Magnesian Limestone. Conglomerate, sandstone, and red marl.
OLD RED SANDSTONE or DEVONIAN.		{ Coal-Measures. Millstone grit. Upper limestone shale and carboniferous limestone. Lower limestone shale ; calciferous sandstones of Scotland.
SILURIAN.	Upper.	{ Upper. Middle. Lower.
	Lower.	{ Ludlow beds. Wenlock beds. Upper Llandovery rocks. Lower Llandovery rocks. Carádac or Bala beds. Llandeilo flags. Lingula flags.
CAMBRIAN.		{ Cambrian sandstones, grits, shales, and conglomerates.

Fundamental or Laurentian Gneiss.

If these various groups and formations had been deposited above each other in regular succession, their united mass would have reached a thickness perhaps twice as great as the height of the loftiest mountain on the surface of the earth. This of course was physically impossible; for if the relative levels of sea and land had remained the same, the land would in time have been entirely worn away, the inequalities in the sea-bed would have been filled up with sediment, and one shoreless ocean would have tumbled round the globe. No more sediment could have been brought to augment the quantity of that which had been already deposited, and hence the total depth of stratified rocks which could have been formed must have been comparatively limited. Instead of an unbroken sequence, however, the stratified portion of

the earth's crust presents numerous proofs of interruption. We see how at one time a certain region extended as a wide surface of land; how in after ages it sank beneath the sea, and became covered over with a thick series of marine deposits; how again it rose and received new races of plants and animals, which lived and died on the soil that had gathered over the hardened sand and mud of the old sea-bottom, wherein shells and corals lay entombed by thousands; how perhaps the same district once more settled down beneath the sea, while a new succession of layers of sand and mud, many thousands of feet in thickness, gathered over its submerged forests, until finally it reappeared as dry land, and received its present vegetation and inhabitants. By this means, while the existing proportions in the distribution of sea and land may not have varied much in past time, there has been, nevertheless, a constant interchange of land and sea over the surface of the globe, one part sinking beneath the ocean, another rising above it. Although, therefore, one unbroken deposition of sedimentary rocks has not been going on continually over the whole surface of the globe, there has yet been always a deposition in progress somewhere. It is the task of the geologist to piece together such fragmentary remains of these scattered strata as may have been preserved, to gather up the evidence which they may furnish as to the progression of life and of terrestrial change, and thus to arrive at some conception of the past history of our planet.

It has just been said that the stratified rocks afford abundant proofs of interrupted succession—that is, they show that a group of strata which was in course of formation at the sea-bottom was upheaved, exposed to denudation, worn down to a greater or less extent, and at last covered over with another later series of deposits. This upheaving of rocks is an important element in the geological economy of nature. It points to the existence of an expansive force lodged within the crust of the earth, manifesting itself sometimes in the form of slow movements of uprise or depression, sometimes in sharp earthquake shocks, and sometimes in the action of volcanoes. These subterranean movements are the origin of the second or IGNEOUS group of rocks, which include lava, pumice, scorïæ, and the other products of an active volcano, together with a large number of rocks found in the crust of the earth, and whose existence there is to be traced to the ejection of mineral matter from some highly-heated source below.

METAMORPHIC rocks are strata which have been subjected to a process of change within the crust of the earth, whereby they have been altered from the condition of sandstones, shales, and other ordinary sedimentary deposits, into hard crystalline masses, in which the signs of stratification are more or less effaced. The cause of this change or *metamorphism* is not yet well understood. It can be traced in all stages of its progress from common sandstone and shale into greywacké and slate, thence into various *schists*, as *mica-schist*, *talc-schist*, *chlorite-schist*, according to the mineral composition of the rock, thence into *gneiss*, until at last we reach granite, the most crystalline and amorphous of the whole, and apparently the ultimate stage of the process. Whatever may have been the origin and mode of this change, it probably went on far below the surface; and in some cases, though possibly not in all, it was accompanied by such a degree of heat as fused the rocks below, and injected the molten mass into

every crack and fissure into which it could find its way. Hence in some of their relations the metamorphic are closely connected with the igneous rocks. Heated water, charged with alkaline or other solutions, has perhaps had a large share in the process of metamorphism.

Each of these three classes of rocks is abundantly developed in the British Islands.* The stratified class covers, of course, by far the largest extent of surface. The igneous group is chiefly developed along the great central valley of Scotland, and along the west coast northwards from the basaltic plateau of Antrim. But igneous rocks occur also abundantly in other districts, as shown on the Map. They appear to be wholly absent, however, from the south-eastern half of England. A line drawn from the Yorkshire coast about Scarborough to the mouth of the Severn divides the kingdom into two nearly equal parts, whereof that to the north-west contains numerous local protrusions of igneous rocks, while that to the south-east shows none. The metamorphic series finds its great development in the Highlands of Scotland, but it also occupies a considerable space in the north of Ireland. Metamorphism is likewise seen round the granite bosses in the Lower Silurian hills of the south of Scotland, among the Lower Silurian rocks of Anglesea, among the Devonian rocks of Cornwall and Devon, and in the Carboniferous series round the granite of Dartmoor.

By comparing this Map with an ordinary geographical one of the same region, it will be seen that certain formations occur in the mountainous and hilly districts, while others are confined to the lowlands and plains. A little attention to this subject will soon show that there is a close connection between the external form of the ground and the nature of the rocks underneath the surface. As a general rule, the high parts of the country consist of hard rocks belonging to the oldest or *palæozoic* division, while the low grounds lie usually upon soft strata of the later or *secondary* and *tertiary* groups. The Highlands of Scotland, for example, are made up of hard tough crystalline rocks, such as gneiss, mica-schist, quartz-rock, granite, and porphyry. The mountains of Wales and Cumberland consist of compact Silurian sand-stones and grits, with thick beds of old volcanic lavas and tuffs. In Ireland, too, the Wicklow chain of mountains is one of granitic rocks. The great plains of central and south-eastern England, on the other hand, lie upon the comparatively soft sandstones, marls, and clays of the new red sandstone, oolitic, and eocene formations.

**PLATES VI. to X.—The Mountains and Table-Lands of Europe, Asia, Africa,
and America.**

This series of Maps is intended to convey correct information regarding the *relief* of the continents. The *physical* position of a place, or its elevation in the atmosphere above the level of the sea, is an element of as much importance in regard to its climate, health, vegetation, and other circumstances, as its *geographical* position, or its vicinity to, or distance from, the Equator. This information on the physical or natural

* It has not been deemed desirable to describe here the distribution of the various formations across the country, as this is sufficiently indicated upon the Map. The nature of the formations themselves, with their characteristic fossils, may be learned from a geological text-book.

geography of country cannot be conveyed in an ordinary geographical map, which must necessarily contain many names and political divisions, to the exclusion or obliteration of purely physical features. The forms of relief on the surface of the globe are varied to an almost infinite extent, yet they may be classified according to their most prominent characteristics.*

These are—1. *Mountains*, which, disposed in linear chains or scattered in isolated groups, extend over the surface of plains and table-lands in every variety of form.

2. *Table-lands* or *plateaus*—elevations of the surface rising to a considerable height, and presenting the form of a table or platform; and—

3. *Plains* and *valleys*, or *lowlands*—portions of the surface little or not at all raised above the level of the ocean.

On comparing the great mountain-chains of the different continents, the first observation which occurs is, that in the Old World—Europe, Asia, and Africa—they follow, in general, the direction of the parallels of latitude, or extend nearly in a direction from east to west; while in the New World, North and South America, the principal chains extend in the direction of the meridians, or from north to south. And next, that in all cases the table-lands are intimately connected with the mountain-ranges—the highest mountains invariably rising, not from plains, but from table-lands. The Old World is most remarkable for its mountains and table-lands, and the New World for its plains.

EUROPE, a peninsular prolongation of Asia, of which it forms the western portion, presents a surface varying in elevation from the plains of the Netherlands, at or below the level of the sea, to the Alps, nearly 16,000 feet above that level. The great mass of the Continent is divided into a region of mountains and high lands in the south-west, and one of low lands and plains in the north and east. The mountainous girdle extends from the Atlantic eastward to the Black Sea, attaining its greatest elevation in *Mont Blanc*, 15,781 feet above the sea-level, the highest point of the *Alps* and of Western Europe.† The Alps constitute the centre of High Europe, and form a network of wild and broken ranges, in which are enclosed narrow but fertile valleys. The tops of the mountains are covered with perennial snow, and from them glaciers descend to the plains. The *Apennines* stretch from the southern part of the Alps south-east throughout the entire length of Italy, forming a long narrow chain from which wide valleys depend. The highest point of the range is the *Gran Sasso d'Italia*, 9543 feet above the sea. The *Fulian* and *Dinaric Alps* unite the *Apennines* with the *Balkan* in Turkey and the *Pindus* range in Greece. The *Carpathians* extend from the Eastern Alps in a semicircular form, enclosing the plain of Hungary on the north and east, terminating in the Transylvanian Mountains in the south. The *Tatra* group in the north rises to 8685 feet above the sea. South-west of the Alps the *Pyrences*, in which *Pic Nethou* is 11,168 feet above the sea, form a natural barrier between France and Spain, and are continued westward to the Bay of

* The main features of the hydrography of these continents are noticed in the paragraphs which accompany Plate XII.—River Systems.

† On the south-eastern verge of Europe, within the political boundary of the Continent, two mountain-peaks—*Elbruz* (18,572) and *Kasbek* (16,545)—the chief summits of the Caucasus range, are respectively much higher than the culminating point of the Alps, but *Mont Blanc* is the monarch among the mountains of historical Europe.

Biscay by the somewhat lower range of the Cantabrian Mountains. The other mountains of the Spanish peninsula are the sierras of Guadarrama and Estrella, near the centre; Morena, and Nevada, towards the south; the last is the most elevated range in the peninsula, its summit, Mount Mulahacem, being 11,678 feet above the sea.

The secondary mountains of Central Europe are the *Cevennes*, in the south-east of France; the mountains of *Auvergne*, with the volcanic group of *Cantal* near the centre; the *Jura* Mountains, between France and Switzerland on the north-east; the *Vosges* on the west and the *Schwarzwald* on the east of the Rhine; the little *Carpathians*, between the rivers Waag and March, in the east; the Harz, of which the Brocken, 3740 feet, is the highest point; the *Erz Gebirge* or Ore Mountains, between the Elbe and Eger, and the *Riesen Gebirge* or Giant Mountains, in which Schneekoppe (the Snowy Peak) rises to 5254 feet, and the low ranges of the *Hundsruok* and the *Taunus* on the north-east and south-west of the Rhine, in Germany. Separated from these central groups by the great plain of France, the Netherlands, Denmark, and the North Sea, are the Mountains of Scotland, the culminating point of which, and of all Britain, is Ben Nevis, in the Grampians, 4406 feet high; and Wales, in which Snowden, the highest point in South Britain, is 3590 feet above the sea; the Alps of Scandinavia, forming a huge barrier of plateaus and mountain-ranges fronting the Atlantic on the north-west and the Ural Mountains on the east. The portions of mountains covered by perennial snow in Europe are limited to the more elevated regions of the Alps, the Scandinavian Mountains, the Pyrenees, and the Sierra Nevada. They are indicated by being left white on the brown ground. These mark the highest points of a system, the sites of true glaciers, and the sources of important rivers.

The TABLE-LANDS are, in Spain, the *Plateau of Old Castile and Leon* in the north, and that of *New Castile and Estremadura* in the south: the northernmost of these is the largest and most elevated in Europe, its average height being 2700, and its highest point 4500 feet above the sea. The *Plateau of Bavaria* forms a terrace on the north side of the Alps, divided into two parts by the Lake of Constance. The only others are the *Plateau of Auvergne*, with *Cantal*, in France; and that of *Bohemia*, an elevated basin, 960 feet above the sea, nearly surrounded by mountains.

The great PLAINS of northern and eastern Europe extend from the Atlantic to the Ural Mountains, and from the Caucasus to the Arctic Ocean. So uniform is the surface here that the entire space may be traversed without changing the level more than a few hundred feet. The Valdai Hills, near the sources of the Volga, 1100 feet, are the highest parts in the interior.

The *Lowlands of France* on the west merge into the *Germanic Plain*, including the Netherlands and the plain of Denmark; and this is extended eastwards through Russia by the Sarmatian Plain in the centre, and the plains of the White Sea on the north, and those of the Black Sea and the Caspian on the south.

The SECONDARY PLAINS are, in Spain, the basin of the *Ebro* in the north-east, and that of the *Guadalquivir* in the south-west; in France, the valley of the *Lower Rhine*; in Italy, the *Plain of Lombardy* and the *Pontine Marshes*; in Germany, the *Valley* of the Lower Danube, the *Hungarian* and the *Wallachian* plains; and in

Scandinavia, the eastern shores of Sweden sloping to the Baltic and the Gulf of Bothnia. In Great Britain are the *Central Plain*, the valley of the *Severn*, and *Plain of York*, in England; and *Strathmore* and the *Carse of Falkirk* in Scotland. In Ireland, the *Great Plain* occupies the greater part of the interior of the island.

The general distribution of the CLIMATE of Europe is indicated on the map by the annual isotherm of 41° Fahr., and the monthly isotherm for January, and by lines tracing the limits of some of the more characteristic trees, grains, and fruits.

ASIA is the largest of the continents or great divisions of the globe, comprehending all the countries of the Eastern Hemisphere east of Europe and Northern Africa. A central mass of continent forms nearly four-fifths of the entire area, and from this extensive peninsulas project on the east, south, and west. Like Europe, it is divided into a vast elevated mountainous region, and a region of valleys or plains. 1. The table-land of Central Asia forms an upheaved region in the middle of the continent, and is crowned by the great wilderness called Gobi or Shamo; its southern border extends from the Indus to the eastern shores of China, supported by the Himalayas, which are prolonged eastwards by the Chinese mountains south of the Yang-tse-Kiang. The broken eastern descent of the table-land extends from the Yang-tse-Kiang on the south to the Amoor on the north; the northern border is girdled by ranges of mountains, of which the Altai is the nucleus; these ranges meeting the Himalaya in the west, form the high land of Turkestan, on which rises the *Bolor Tagh*, called by the natives the *Roof of the World*, the crests of which are supposed to be 19,000 feet above the sea. This table-land is crossed in an easterly and westerly direction—first, by the *Thian Shan* or Celestial Mountains, on which is the high peak of *Peshan*; second, by the *Kuenlun*, which is extended eastward by the *Pe-ling* Mountains; and between these is the plateau or high valley of Eastern Turkestan. Between the Kuen-lun and the Himalaya proper is the *Table-land of Tibet*, which appears to be composed of range after range of snow-clad mountains, connected with, and not much inferior in elevation to, the Himalayas.

The mountain systems of the *Altai* and its offsets separate Mongolia on the south from Siberia on the north. Extending north-east from the Altai, the mountains of Dauria are prolonged by the Yablonoi Mountains to the Sea of Okhotsk, and the Stanovoi and Aldan Mountains stretch north-west and north-east to the arctic circle. The *Inshan* and *Khingan* ranges form the south-eastern buttress of the plateau of Mongolia.

The table-land of *Iran* or *Persia* is connected with that of Central Asia by the Hindoo Koosh Mountains: it has an average height of 2500 to 3500 feet, and on its north side the Volcano of Demavend in the Elburz range is 18,464 feet in elevation. This is joined on the west by the plateau of *Armenia*, about 7000 feet high, supporting Mount Ararat, on which the ark is supposed to have rested after the Flood, 16,964 feet above the sea; and that of *Asia Minor*, a mountainous mass with a mean elevation of about 3280 feet, on the south side of which are the Taurus Mountains, 9800 feet, and Mount Arjish, 12,600 feet above the sea. The *Table-land of Arabia* appears to have a general elevation of 2000 to 4000 feet; on the south-east, the mountains are 5000 feet above the sea-level: a large portion of the interior is an arid desert. The *Mountains of Syria and Palestine* form a connecting link between the table-land of Arabia

and the Taurus ; the highest point of Lebanon has an elevation of 10,061 feet above the sea.

On the north the peninsula of *Hindustan* is bounded by the *Himalayas* ; and on the south, the *Western and Eastern Ghauts*, running parallel with the shores, enclose the table-land of the Deccan, which is bounded northwards by the Vindhya Mountains. The average elevation of the Himalaya is probably 16,000 to 18,000 feet. The highest peaks are Dhawalagiri, 26,826 feet ; Kanchinjanja, 28,156 feet ; and Mount Everest, situated nearly midway between them, 29,002 feet, or $5\frac{1}{2}$ miles, above the sea, the highest known point on the surface of the globe. The table-land of the Deccan has a mean elevation of 1500 to 2500 feet above the sea, and is covered with isolated conical hills rising 2000 feet above the plateau. The *Mahableshwar Hills* in the Western Ghauts are 4700 feet above the level of the sea : southward in the *Neilgherry Hills*, the highest point of the Ghauts is reached in Dodabetta, 8640 feet. In Ceylon, Pedrotallagalla is 8280 and Adams' Peak, 7420 feet above the sea.

The *Plains* of Asia comprise a large portion of true steppes and deserts, interspersed with fertile valleys. The *Chinese Lowland*, a district of pure alluvium, extends from the shores of the Yellow Sea to a great distance inland, and occupies the deltas of Yang-tse-Kiang and the Hoang-ho. The plains of *Further India* include the rich valleys on the west of the Gulf of Tonquin, and those of the rivers Mekong, Meinam, Saluen, and Irrawady. The *Plains of Hindostan* comprise, on the east, the valley of the Ganges, which, at its mouth in the Bay of Bengal, forms a cluster of marshy islands called the Sundarbans. On the west, the *Valley of the Indus*, an alluvial plain, extends for many hundred miles without any eminence, except the Salt range of hills near the western base of the Himalaya. In the south it is mostly a desert, called the Thar, and it contains the remarkable depression of the Runn of Cutch. The lowland of *Turan and Bucharia* extends from the table-lands of Iran on the south to the Ural Mountains and a western branch of the Altai on the north ; its lower portion contains the great depression of the Caspian and Aral seas, forming the central part of a great basin of *Continental drainage*, the largest hollow on the land surface of the globe. The *Plains and Steppes of Siberia* extend from the Ural Mountains on the west to Behring Strait on the east, sloping northward from the base of the Altai and the Stanovoi mountains on the south to the Arctic Ocean on the north. In the southern regions the surface is covered with forests, and has fertile valleys ; in the west the steppes are bare, flat, and undulating ; and in the north, bordering on the Frozen Ocean, it is a trackless wilderness, with numerous salt lakes and marshy flats, called *Tundras*, the soil of which is permanently frozen to a depth of several hundred feet.

The *Lowlands of Syria and Arabia* are in the south dry and stony deserts, but the lower courses of the Tigris and Euphrates in the north are rich and fertile. In the south of Palestine, the Dead Sea is the deepest and most remarkable depression on the face of the globe, its surface being 1298 feet below the Mediterranean.

AFRICA, the south-western extension of the mainland of the Old World, may be regarded as an immense peninsula of the European and Asiatic continent, separated from it naturally by the Mediterranean and Red Seas, and artificially by the Canal of Suez, which has been cut through the narrow connecting isthmus.

Unlike the chief continent, in which the mountains and highlands are grouped near the centre and ramify outwards to form long peninsulas, with deep bays between, the main features of Africa resolve themselves into one vast highland plateau in the east and south, and a great lower region in the north-west; the land is compact in form, the shores are rounded and deficient even in river entrances, and the coast islands are few.

The whole of Africa south of the 10th parallel of N. latitude, so far as it is known, is a continuous highland, rivalling that of Asia in extent, though not in elevation. On the north-east it runs out to form the vast wedge-shaped plateau of Abyssinia, which is again prolonged by lesser elevations as far as the Isthmus of Suez. The eastward side of the African highland is the highest part of the continent, and its fall towards the Indian Ocean is abrupt and steep, whilst the slope is more gradual towards the Atlantic. Some of the highest mountains of the continent stand close to the eastern verge of the plateau. From Jebel Attaka, immediately west of Suez (2640 feet), the rise of the eastern side begins. About midway in the African coast of the Red Sea, Mounts Elba and Soturba each attain an elevation of over 6000 feet. Near the apex of the Abyssinian wedge Mount Abba Jared rises to 15,000 feet, and Ras Detschen to 15,986 feet, and the whole line of the plateau edge facing the lowland which skirts the Red Sea is higher than 7000 feet. Towards the equator, still on the verge of the highland, are Mounts Kenia (18,000) and Kilimanjaro (18,700 feet), each snow-clad at its summit, and in their neighbourhood stand many less known but possibly higher peaks or "Doenyos," as the isolated summits in this region are named. The active volcano called the Njemsi Mountain has frequently been reported in this region. Continuing southward, the eastern edge becomes the Rubeho Mountains, a range which is crossed by the travellers from Zanzibar towards the lake region, and is known again where it walls in the great lake Nyassa as the range of the Njesa Mountains. Nearer the Zambezi it is marked by Mount Zomba, 7000 feet; Mount Milanje, 8000 feet; and Mount Clarendon, 6000 feet above the sea. The valleys of the Zambezi and Limpopo rivers form the only considerable break in the continuity of the edge, but between these it is still marked by the mountains of the South African gold region, where Mount Matoppos is 7200 feet in elevation. The plateau again approaches more nearly to the coast in the cliffs of the Drakenberg or Quathlamba, on the slopes of which lies the colony of Natal. Cathkin Peak, near the middle of these heights, is 10,380 feet above the sea. The inmost and highest ridge of terraces which form the Cape Colony is the direct continuation of the Quathlamba, and Compass Berg, at the south-eastern curve, is 8500 feet high. Westward this inner edge decreases in elevation in the parts of it which are known as the Nieuweveld and Roggeveld. The crests of the two outer and lower terraces pass round the Cape Colony in lines parallel to the coast. Table Mountain, a well-known flat-topped height (3550 feet), is an isolated granite mass, overhanging Cape Town, and forming the nucleus of the peninsula which extends to the Cape of Good Hope.

The western side of the South African plateau is generally much lower than the eastern, is not so abrupt in its descent, and is broken through by many rivers, the chief of which are the Orange, Cunene, Coanza, Congo, and Ogowai (pronounced

Ogōwee), but it still forms a well-marked edge, generally parallel to the coast-line. Among the more prominent portions of this side are the mountains of Namaqua and Damara Lands; Mount Omatako, in the latter region, attains 8800 feet. The more broken serras and terraces of Benguela and Angola are also of considerable elevation; rapids and cataracts mark the passage of the rivers from the plateau to the lower land bordering the sea. Towards the equator the edge is again more distinctly marked, and, near the Ogowai River, Mount N'Kumu Nabuali is reported to have an elevation of upwards of 12,000 feet. Several active volcanic mountains of great elevation are also reported in this neighbourhood, though these have not yet been seen by any European. The terminal point of the edge, on this side of the continent, is marked by the volcanic knot of the Cameroon Mountains, 13,700 feet high, rising in a peninsula nearly detached from the north-western corner of the plateau.

The north-facing side of the vast plateau is the least known, but is believed, nevertheless, to form a distinct and perhaps prominent line, since the rivers, flowing from its supposed place, have the character of great mountain streams, rising and falling rapidly at different seasons. Mounts Alantika, 9000 feet, and Mindif, 6000 feet, discovered by the traveller Barth, south of Lake Chad, appear to be outliers of the plateau. Eastward this edge is shown in the heights which narrow the Nile Valley, and cause the cataracts above Gondokoro, and, further still, it joins into the inner slope of the Abyssinian plateau, turning with it to northward.

Among the more prominent elevations of the South African plateau, within the limits just described, are the Blue Mountains, which rise on the western side of the Albert N'yanza to an elevation of about 10,000 feet; these are believed to be continued southward to near the third parallel of S. latitude, in the great highland of Ulegga, mentioned by Speke, Baker, and Livingstone, which again joins into the known mountains which separate the head of the Tanganyika from the Albert Lake, and which rise in Mount Mfumbiro to 10,000 feet. This line of heights, taken in its connection with the high plateau land of Unyamuezi, south of the Victoria Lake, is the most important water-parting of the continent, since it certainly limits the Nile basin in the south, and contains its long-sought sources. The ridge which divides between the Tanganyika and Nyassa Lakes, called by Livingstone the Lobisa plateau, extends also round the head lakes of the Chambeze (or Congo). In this part it was named the Serra Muchinga or Maxenga by the Portuguese travellers who first crossed it, and its extension north-westward forms the Kone Mountains of Livingstone. Almost in the same latitude the ramifying knot of the Mossamba Mountains has the sources of rivers flowing to all parts of the compass. Elsewhere the plateau has a more uniform surface, of a general altitude perhaps not exceeding 3000 feet.

Beyond the northward slope of the great southern plateau, almost the whole of the remaining portion of Africa stretches in a great lower land, of a general elevation of perhaps less than 1000 feet to the Mediterranean. On the south-west it has a higher margin, formed by the mountains of Kong, which rise from 2000 to 3000 feet, and extend parallel to the coast-line from near the mouth of the Niger to the Gambia, and supply the main tributaries of the former river. On the north-east, also, there is a highland named the plateau of Barbary, which extends through Marocco, Algeria,

and Tunis, enclosed between a double range of mountains. In its western portion, where the outer ranges draw together, the elevation of the plateau of Barbary is very considerable, and its summit, Jebel Miltsin, attains the height of 11,400 feet. By far the greater part of the northern lower land is occupied by the vast region of the Sahara, characterised by great belts of sand-hills, and by wide stony plains; though scattered between these are higher mountain-knots with rich vegetation, and deep "wadis," or "oases,"—the beds of periodical streams which carry with them lines of the highest fertility. Some of the most important of these wadis descend from the inner slopes of the plateau of Barbary.

Two areas of depression beneath the sea-level are known in the African continent, and form a remarkable feature in its relief. The most considerable of these is in the Lybian desert, extending from near the lower valley of the Nile to the great Syrtis westward, only separated from the Mediterranean by the belt of high ground of which the small plateau of Barca is the chief elevation; the interior extent of this depression is unknown, but it may well be compared with that of the Caspian Sea in Asia, and is in some parts from 100 to 150 feet beneath the sea-level. The triangular space enclosed between the steep eastern slope of the plateau of Abyssinia and the Red Sea coast is also an area of depression, and includes a number of salt lakes, one of which (Lake Assal) is believed to be not less than 600 feet below the Red Sea. Mount Artali, in this depressed region, is one of the few known active volcanoes of Africa.

Africa is the only one of the continents of the globe which has a large extent of land on each side of the equator, and upon its surface, therefore, are marked more clearly than in any other part of the world those natural landscape belts which result from the unequal distribution of heat and moisture in different latitudes. The central portion, extending for several degrees on each side of the equator, in the region of frequent tropical rains and great heat, is characterised by dense forests, intertwined with the most luxuriant and rank vegetation. To north and south of this are corresponding belts of less wooded park-like country, merging into pasture-land and almost treeless grass countries. These pasture-belts gradually change as the dry, almost rainless, regions of the Sahara in the north, and of the Kalahari desert in the south, are approached, and in these, sandy and stony waste lands are characteristic. Beyond the deserts again, where the moisture from the Mediterranean, and from the Atlantic southward, is carried inland, the coast slopes of the plateau of Barbary and those of the Cape colonies, corresponding, are fertile lands, either pastoral or cultivated.

NORTH AMERICA comprises all that portion of the American continent north of the Isthmus of Panama; its area is more than twice that of Europe, and more than one-sixth of all the dry land of the globe. Its axis of elevation is on its western side, whence its surface slopes to the east, north, and south-east. A vast series of table-lands, more than 3000 feet in mean elevation above the sea, forms a broad irregular belt on the Pacific side, extending from Behring Strait to the Caribbean Sea. Along the western edge of this plateau, from the territory of Alaska to the peninsula of Lower California, stretches the *Oceanic Mountain chain*, parallel to the Pacific. At its northern extremity are the volcanic peaks of *St Elias* and *Fairweather*, rising to

nearly 15,000 feet. South of this the *Sea Alps* are succeeded by the *Cascade Range*, where *Mount Baker* is 11,000 or 12,000 feet, *Mount St Helens* 9750, *Mount Hood* 11,225, and *Mount Shasta Butte* 14,440 feet above the sea. Between the parallels of 35° and 40° the *Coast Range* and the *Sierra Nevada* enclose the fertile gold-producing valley of California in the basins of the Sacramento and Joaquin rivers. Several peaks of the Sierra Nevada attain a height of 10,000 feet. *Mount Dana* is 13,000 feet above the sea. South of this the mountains are prolonged through the Sierra de Santa Lucia (in which Mount Whitney is 15,000 feet high) to the peninsula of Lower California.

The *Rocky Mountain chain*, the eastern side of the plateau, forms the principal watershed of the continent, all the waters west of this flowing to the Pacific, and all east to the Arctic and Atlantic Oceans. It comprises two, and in some places three, distinct ranges of high mountain-crests, table-lands, and valleys. The main chain forms a waving line curving along the eastern part of the table-land; it is not continuous, but broken by deep passes, and the leading ridges often overlap each other. Its easternmost range contains mountain-peaks rising 16,700 feet above the sea. North of the River Platte, the *Wind River Mountains*, the highest of the chain, rise in *Fremont's Peak* to 13,750 feet. This lofty axis separates the head-waters of the Missouri, flowing to the Atlantic, from those of the Columbia, which flow westward to the Pacific Ocean. North of the Wind River chain, the eastern range, near the sources of the Saskatchewan, rises in Mount Hooker to 16,750 feet, Mount Murchison to 15,789 feet, and Mount Brown to 16,000 feet above the sea. From this point the mountains of the eastern range decline northwards to the Arctic Ocean. South of the Wind River Mountains the eastern ridge is marked by *Union Peak*; *Long's Peak* 14,056 feet; and *James's Peak* and *Pike's Peak* 14,216 feet; and continued southward to the *Cordillera of Coahuila* and *Potosi*, forming the eastern edge of the Mexican table-land. Its western ridges are the *Sierras Verde*, *Mimbres*, and *Madre*, enclosing on this side the *Table-land of Mexico* or *Anahuac*, which has an elevation of from 6000 to 9000 feet above the sea. The plateau is in some places traversed by well-defined ridges, but in others it is unbroken either by hills or depressions. The City of Mexico is 7460 feet above the sea, and on the south-eastern and loftiest side of the table-land the volcano of *Popocatepetl*, the highest point of the Mexican system, is 17,783 feet high. East of this is the volcano of *Orizaba*, 17,879 feet above the sea. The *Volcanic Range of Guatemala* comprises a great number of volcanic peaks, rising from a table-land about 5000 feet above the sea: of these, *Agua*, 13,200 feet in elevation, is verdant to the summit. Immediately west of it the volcano of Fuego is 13,000 feet high.

Between the Rocky Mountains on the east and the coast-chain and Sierra Nevada on the west there extends, from the Gulf of California to the Arctic Ocean, a high desert zone, which in its central regions, between lat. 35° and 45° N. in the *Great Basin of Utah*, is from 4000 to 5000 feet above the sea. This region is nearly rainless, and comprises wide plains encrusted with salt. It contains twelve or fifteen saline lakes, the largest of which is the Great Salt Lake of Utah, 4210 feet above the sea.

The *Mountains of Veragua* are upwards of 5000 feet above the sea level, and the

Isthmus of Panama is about 850 feet in mean elevation, but the Pacific Railway crosses at a height of only 262 feet above the sea.

The *Appalachian* or *Atlantic Mountain System* (Alleghanies), on the east, is the only counterpoise to the vast mountain-mass on the west side of the continent. It embraces the entire range of high lands extending from the Gulf of St Lawrence to Georgia, about 2000 miles in length, and 150 to 200 miles in breadth. It is composed of many parallel ridges, with crests reaching to 6700 feet, their bases resting on a plateau, the mean elevation of which is 2500 feet above the sea. At the south end of the Blue Ridge, Black Dome is 6707 feet high.

The *Great Central Plain* of North America extends northward from the Gulf of Mexico to the Arctic Ocean, between the Rocky Mountains and the Appalachian chain. It is a vast region of plains, slopes, and table-lands, watered by many great rivers, and indented by large lakes; on the west it forms a region of dry elevated steppes or table-lands, and on the east a wider and lower tract of hills and rolling plains generally well watered. Near the middle of the plain a broad gentle swell of the land crosses the continent from the coast of Labrador, north of Lake Superior, west-south-east towards the Pacific coast. In the *Missabay Heights*, west of Lake Superior, it is 1500 feet above the sea, and divides the waters flowing north to Hudson Bay, from those which flow south to the Gulf of Mexico.

The *Atlantic Slope* stretches from the Gulf of St Lawrence to the Gulf of Mexico, narrow in the north, but expanding southwards to 200 miles in width from Virginia to the southern end of the Alleghanies; its greatest height is about 1000 feet, near the sources of the Roanoke. The *Atlantic Plain*, or seaboard belt of the Atlantic slope, is nowhere more than 100 feet above the level of the sea; in the south-west its seaward portion is swampy and overflowed. This lower plain continues along the shores of the Gulf of Mexico, the peninsula of Yucatan, and the shores of Honduras, terminating in the valley of the San Juan and the Lake of Nicaragua.

The continent of North America comprises a broad *forest zone* extending from Alaska in the north-west to the Alleghanies in the south-east, a narrow zone of forests on the Pacific coast, and an almost *treeless region* extending from the Pacific-coast mountains to the Mississippi. The northern limit of trees cuts the north of Labrador, and passes north of Great Slave Lake. North of this is the *Arctic treeless region*. On the map are marked the northern limits of the growth of wheat, and of the wine-yielding vine, as well as the annual isothermals of 40°, 50°, and 60° Fahrenheit.

In **SOUTH AMERICA**, as in the northern portion of the continent, the elevated axis is on the west or Pacific side, where the great mountain-system of the Andes stretches, through 65° of latitude, from Cape Horn to the Isthmus of Panama, with a breadth of 40 to 400 miles. The centre of the continent is occupied by the immense plains of the Amazon, the Plata, and the Orinoco; and in the east are the separate mountain-systems of Parimé and those of Brazil. The *Cordillera* or *Great Chain of the Andes* is divided, according to the countries which it traverses, into five principal sections or groups: (1.) The *Andes of Quito*, extending from lat. 5° S. to the Isthmus of Panama in lat. 8° N. In this-division more of the giant mountains are congregated than in any other part of the chain. On the east side the snow-clad summit of

Antisana is 15,137, and the beautiful white cone of *Cayambe*, traversed by the equator, 15,532 feet above the sea, and on the west *Pichincha* is 15,924 feet in elevation. Between these ranges lies the table-land of *Quito*, with the city of *Quito*, 9,520 feet above the Pacific. In the chain of *Quindiu* is the peak of *Tolima*, 18,130 feet, the highest point of the Andes north of the equator. (2.) The *Andes of Peru* extend from lat. 15° to 5° S. Near the south-east extremity is the mountain-knot of *Vilcanota*, 17,525 feet. In about lat. 10° is the mountain-knot of *Huamanga* and *Pasco*, 11,500 feet, on which rises the *Nevada de la Fria*, 16,000 feet. (3.) The *Andes of Bolivia*, the central and most elevated portion of the system between lat. 21° and 15° S., consist of two great parallel mountain-chains—the *Cordillera* of the coast, and the eastern or *Bolivian Cordillera*—enclosing a vast Alpine valley, the plateau of Bolivia, partly occupied by the lake of *Titicaca*, 12,347 feet above the sea. In the western chain are the *Nevada of Gallinieri*, 21,900 feet; *Sabana*, 23,014 feet; the volcano of *Arequipa*, a regular cone, 20,320 feet; and the *Nevada of Copacabamba*, 21,000 feet above the sea. The eastern chain, or the *Cordillera Real*, has a mean elevation of 16,000 feet. In it rises the *Nevada of Illimani* with an elevation of 21,338 feet, the lowest glaciers on the northern side of which do not descend below 16,500 feet. North-west of this is the *Nevada de Serrata*, or *Aurumani*, the most elevated of all this snow-capped range, perhaps the highest point of the American continent, 24,812 feet. (4.) The *Andes of Chile* extend from lat. 21° to 42° S., running for a considerable extent in a single ridge of about 30 miles in breadth. The mean elevation of the chain is about 12,000 feet, but several serrated peaks rise to a much greater elevation. Visible from the port of *Valparaiso* rises the giant of the Chilean Andes, the peak of *Aconcagua*, 22,421 feet above the sea-level. Inland from this, near the capital of Chile, is the *Nevada of Tupungato*, 20,269 feet. There appear to be five active volcanic vents in the southern prolongation of the Andes of Chile—viz. *Maypu*, *Chillan*, *Antuco*, *Villa Rica*, and *Oserno*. (5.) The *Patagonian Andes* commence at the Strait of Magellan, and run north to lat. 42°, with an elevation of 3000 to 8000 feet above the sea. Their summits are covered with perennial snow, whence glaciers descend almost to the sea-shore. The chief of these snowy mountains are *Osorno*, 7550 feet above the sea; *Yanteles*, 8030; *Melimoyu*, 7400; and *Stokes*, 6400 feet. Under the parallel of lat. 20° S. the snow-line rises to 18,500 feet; in lat. 26° 30' it is 13,800 feet, and on the parallel of *Valdivia* it sinks to 8300 feet.

The *Mountain-System of Parimé* forms a plateau between the *Orinoco*, the *Rio Negro*, and the *Amazon*; the principal chains run in a direction generally from east to west. Near the centre the sierra *Pacaraima* is 7000 feet, and on the west, *Mount Maravaca* is 10,500 feet above the sea-level.

The *Mountains of Brazil* form a vast isolated plateau between the *Amazon*, the *Madeira*, the *Paraguay*, and the *Plata* rivers. In the south the *Serra do Mar* or coast-range, forms the south-east edge of the plateau. This, with the *Serra des Oragos*, or *Organ Mountains* (so named from the resemblance of their peaks to the tubes of an organ), compose the so-called *Brazilian Andes*. Proceeding northward, the other chief ranges are the *Serra do Espinhaço*, *Serra Timba*, *Serra Tabatinga*, and *Serra Irmaos*. The *Cordillera Grande* and the *Serra Santa Martha* divide the waters of the

Uruguay and the Tocantins, and a high ridge south of the Orinoco separates Brazil from Venezuela. The highest points of the Organ Mountains reach an elevation of 7500 feet, and Mount *Itambe* is 5755 feet above the sea-level. South of this the *Serra de Piedade* is 5830 feet, and *Itacolumi* 5750 feet in elevation. Immediately within the coast-range, the *Sertao*, or table-land of Brazil, rises by several gradations to the central plateau, the mean height of which is estimated at from 2500 to 3000 feet.

The PLAINS of SOUTH AMERICA comprise the principal and several subordinate divisions. (1.) The *Llanos* or *Plains* of the *Orinoco*, between the mountains of Brazil on the east and the chain of Suma Paz on the west. These plains are perfectly level, and at a distance of 450 miles from the sea-coast the surface is only 192 feet above the sea. From their peculiar position, encompassed by mountains, within the torrid zone, they are, during one-half the year, desolate sandy wastes, while during the other half, after the rainy season, they are covered with luxuriant grass. (2.) The *Atlantic Slope*, the *Valleys of the Amazon* and of the *Rio de la Plata*. All the large rivers of South America flow eastward to the Atlantic Ocean, indicating the slope of the surface in this direction. The valley of the Amazon, the largest river-basin in the world, is situated partly in the northern, but mostly in the southern hemisphere. Its climate forms a striking contrast to that of the Orinoco, for it is copiously watered at all seasons, supporting a luxuriant vegetation, and its humid plains are occupied by impenetrable forests, the growth of a thousand years.

The *Valley of the Rio de la Plata* consists chiefly of two immense plains, the northern watered by the Salado, Vermejo, Pilcomayo, &c., and the southern comprising nearly all the *pampas of Buenos Ayres*—a vast plain, covered for the greater part with rich pasturage and gigantic thistles, interspersed with saline lakes.

The *Patagonian Steppe* is composed of a series of terraces east of the Andes, covered with marine shells, clay, earth, and gravel: these terrace-slopes are arid and sterile.

The *Pacific Slope* is a narrow belt between the chain of the Andes and the ocean, from 50 to 100 miles in breadth. In the south it comprises the moist shores and islands of Patagonia, in Chile the fertile provinces of Talca, San Fernando, and Concepcion, and from lat. 30° to lat. 10° S. the rainless district of Peru. Its fertile valleys are generally separated by long deserts of sand.

PLATE XI.—Ocean Currents and Cotidal Lines.

I.—OCEAN CURRENTS.

A constant and universal circulation is taking place in the waters of the ocean, and this movement, besides being essential to the preservation of uniformity in the component parts of the sea, is one of the most powerful agents in the distribution of temperature and moisture over the globe. In the most comprehensive sense, this circulation is believed to be due to, and to be sustained by, difference of temperature—the first movement being given by the action of cold near the polar regions of the globe upon the waters in these parts, increasing their specific gravity, and causing them to

descend and flow over the ocean-bed towards the Equator as an under stratum. Such an underflow involves, as its necessary complement, a movement of the upper stratum, bearing with it the warmth of equatorial and temperate seas towards the Poles. This theory appears to be the only one which will account for the presence of glacially cold water in the deepest parts of the ocean, in the temperate and equatorial regions, as observed by the thermometer; and the theory seems to be confirmed by the fact, that in enclosed seas such as the Mediterranean, which have but a narrow and shallow opening to the ocean, and can thus have no communication with the Polar Seas, the temperature of the waters, at depths corresponding to those found in the ocean in the same latitude, is higher, and the deepest waters have a temperature which approximates very closely to that of the crust of the earth in that region.*

Whilst this thermal circulation constitutes the grandest movement of the whole ocean, and takes place at a constant and uniform, though slow, rate of progression, there are a number of more superficial and rapid movements of the sea-surface grafted on to, and merging into, the greater circulation, which form the better known and more apparent currents of the sea. These more sensible, and more or less constant currents, which play an important part in the navigation of the sea, are graphically shown in their general direction and extent upon the accompanying chart. Their presence is due to a number of causes besides that which has been indicated, and which act either separately, or more frequently, in combination one with another. Looking at the chart, it is observed that in each of the great oceans there are westward flowing currents on each side of the Equator. These are believed to have their prime moving force in the rotation of the earth, the inertia of the waters tending to leave them behind in the turning of the earth from west to east; but besides this, they have a powerful aid to their motion in the trade winds. Where a continent intervenes as a barrier to the westward movement of these currents—the waters are pressed against its shores, and tend to escape by the most ready way. The most prominent examples of such escape currents are the *Gulf Stream* of the Atlantic, which is driven with great force out of the Gulf of Mexico by the pressure into that enclosed sea of the Equatorial currents; the *Japan Current* of the Pacific, and the *Mozambique Current* of the Indian Ocean. The two former of these currents are aided after their escape, and carried on to higher latitudes, by the anti-trade or prevailing south-west winds of the regions into which they flow. Once set in motion, these currents give rise to another class of streams which follow to supply the place of the waters which have changed their position, and to maintain the equilibrium of the Ocean surface. Such compensation currents, the peculiarity of which is that they are drawn from in front, not pushed on from behind, are the streams which flow from the Southern Ocean to the Atlantic, the Indian Ocean, and the Pacific, as the *South Atlantic Current* of the West African Coast, the *West Australian Current*, and the *Peruvian Current*, though each of these after its first outset is aided in its movement by the south-easterly trade-wind. The *counter currents*, which are found between the Equatorial streams, may be considered as purely compensational currents, as may also the currents which, moving

* See "Report on Scientific Researches carried on during 1871, in H.M.S. 'Shearwater,'" by Dr Carpenter. —Proceedings of Royal Society, No. 138.

at a slower rate, and generally in the direction of a prevailing wind, connect the more marked and decided ocean streams. The most remarkable examples of currents flowing in diametrically opposite directions side by side are found among these Equatorial streams; the best known is perhaps that of the *Guinea* or counter Equatorial current of the Atlantic, where it approaches the south Equatorial stream in the Gulf of Guinea. The strong current which flows round Cape Horn is perhaps the best example of a constant stream which has its origin in a prevailing wind.

Admitting the probability of the theory that the prime mover of the great oceanic thermal circulation is the action of cold on the waters bordering the Polar circles, causing these to descend to the bottom of the ocean, we should not expect to find any considerable surface outflow of water from the Polar regions; yet the cold surface currents which flow out of both Polar areas, but more especially those from the land-locked Arctic region, have the most important bearing upon the climate of the regions with which they come in contact. These ice-bearing currents, known in the southern region as the *Antarctic drift*, and in the northern hemisphere as the *East Greenland* and the *Labrador streams*, must then be considered as the overflow of water rendered specifically lighter or fresher than the general mass of the ocean by the melting of ice and snow at the margins of the frozen regions surrounding the Poles, their floating power being maintained by the gradual reduction of the icebergs and drift ice which they transport into milder climates. The Labrador current may also be considered as in part a compensation current to the Gulf stream, which flows in a directly opposite direction towards the Arctic basin. Along the line of meeting of cold and warm currents, and where these interlace, fogs most frequently arise; the best known are those of the Newfoundland Bank. *Periodical currents*, due solely to the action of a particular wind, before which they are drifted, are most prominently developed in the monsoon region of the Indian Ocean and the China Seas, and frequently attain great strength where the waters are hemmed in between the East India Islands. Among these barriers their direction is twisted and turned by the form of the coasts, but the general movement is with the wind to northward with the monsoons blowing from the south (from March to September), and to southward before the winds from the north (from September to March).

The rate of motion of the ocean currents is very different in separate parts of the same stream, and the rate at special points may vary considerably at different seasons. The highest observed rates have been found in parts of the currents which are compressed by some peculiar formation of the land which compels their direction; thus a movement of upwards of 100 miles a day has been recorded in the Gulf stream, where it rushes through the narrows of Florida, and in the Mozambique current at Cape Corrientes, but it does not seem probable that any current in the open ocean attains a rate of more than 60 miles a day, and this has only been noticed at exceptional times in the strong equatorial streams; the average rate of the chief currents may, perhaps, be taken at 15 miles a day, whilst some of the drifts which prolong these have a scarcely perceptible movement.

The effect of the ocean circulation in ameliorating climate is nowhere more strikingly exemplified than in the European coasts, where the warm south-westerly

flow from the temperate regions of the Atlantic, in the line of the Gulf stream, keeps the harbours of the British Isles, and of Norway, to its most northerly point, free of ice throughout the year; nor can the adverse power of the polar surface streams be better shown than by contrasting this coast with that of the opposite side of the Atlantic, which is blocked with ice throughout the year, in latitudes south of that of the British Isles.

II.—COTIDAL LINES.

The observed connection between the successive risings of the waters of the sea on all coasts, and the meridian passage of the moon, has established the belief that the moon is the cause of these risings called *tides*. The attraction of this body acts most strongly on those parts of the ocean which are more immediately beneath her, and tends to draw up the waters and raise a tidal wave at each meridian which is successively passed in her course, forming, at the same time, a secondary or lower tidal wave in the opposite hemisphere.

The sun also acts in the same manner, though with less force than the moon, owing to its greater distance, and modifies the form of the ridge produced by the moon's attraction. When the sun and moon are in the meridian together, or in direct opposition, their combined action produces the higher tides, known as *spring* tides, which happen after new and full moon. Again, when sun and moon are 90 degrees apart, relatively to the earth, their attractions tend to neutralise one another, and *neap* tides, or the lowest tides, occur, after the first and third quarters of the moon.

A cotidal line is one drawn through all adjacent parts of the ocean which have high water at the same time, and on the map these have been drawn at intervals of one hour. The line for any hour may be considered as representing the summit or ridge of the tidal wave at that time, or that protuberance of water upon the surface of the ocean, which moves along the seas, and, by its motion, brings high-water and low-water to any point at the time when the elevated and the depressed parts of the watery surface reach that place. The cotidal lines for successive hours represent the changing positions of the summit of this wave. If we suppose a spectator detached from the earth to perceive the summit of the wave, he will see it travelling round the earth, in the open ocean, once in every twenty-four hours, accompanied by another at twelve hours distance from it. If the whole surface of the terrestrial globe were uniformly covered with water, the tides would be governed by the moon and sun only, and high-water at every place in the same latitude would follow the transit of the moon, at the same interval of time, and the wave would travel with very great velocity. The interposition of land continents, the unequal depth of the sea, the presence of arms of the sea, bays, and estuaries, of islands and island groups on the actual surface of the globe, however, interrupt and greatly diminish the movement of the tidal waves, and give rise to the irregular form of the cotidal lines, and to the frequent meetings of the tidal waves from opposite directions, as, for example, in the narrow seas which surround the British Isles. These interrupting lands also give rise to great variations in the velocity with which the tidal wave travels in different parts of the ocean.

Since the ridge of high-water moves from the position of one cotidal line to another in an hour, the velocity of its movement may be estimated from the map: where, as in the Atlantic and Pacific Oceans, the hour lines are perhaps ten degrees of latitude apart, the rate of movement must rise to about 700 miles an hour, whilst, in the English Channel, its progress is limited to about 30 miles. On the other hand, the magnitude of the tide is generally less in the open ocean than in enclosed seas, and the greatest tides and tidal currents occur where waves converge in a channel narrowed by the surrounding land, and where, from the gradually shelving nature of the sea-bed, the tidal wave, advancing through the entire depth of the ocean, is forced gradually upwards by the wedge-shaped surface over which it moves. Such a formation causes the phenomenon known as the *bore*, when, during spring tides, after a low ebb, the tidal wave advances up the sloping and narrowing passage with great rapidity. Remarkable examples of this phenomenon are known in the Runn of Cutch in India, in the Bay of Fundy, and in the mouth of the Amazons in America; nearer home in the Solway Firth and the Bristol Channel.*

PLATE XII.—River Systems.

In this map, the land surface of the globe has been marked out and coloured to show the great river systems which belong to each chief division of the ocean, and to indicate the great areas of continental or interior drainage, from which no water returns directly to the sea. Thus the pink colour shows the area of land, the rivers of which flow to the Atlantic; the blue the drainage to the Arctic Sea; the yellow to the Pacific; and the green to the Indian Ocean. Areas of continental drainage between these are coloured purple.

Since the great mountain regions of the world which form the chief water-partings—such as the Rocky Mountains and the Andes in America, the mountains of China, of India and Persia, and of eastern Africa—rise uniformly nearer to the coasts of the Pacific and Indian Oceans than to those of the other seas, it follows that the longer slopes and the great plains of the continents lie towards the opposite side. Hence it is that the area of drainage to the Atlantic and Arctic Seas, as shown in the following table, is double that of the steeper slopes which fall to the Pacific and Indian Oceans, and that almost every one of the great rivers and fresh-water lakes of the world are found in the former systems. To these belong the *Mackenzie* River, the basin of which forms the chief part of the Arctic drainage of America; the *Nelson* to Hudson Bay; the *St Lawrence*, *Mississippi*, *Amazons*, and *Parana* flowing more directly to the Atlantic. In the old world, belonging to the same systems, are the *Congo*, rising in the newly-discovered lake chain near the eastern side of Africa; the *Niger*; the *Nile*, from the equatorial African lakes; the *Danube* and *Rhine* in Europe; and the great rivers of Siberia forming the Arctic drainage of Asia, the *Obi*, *Yenisei*, *Lena*, *Indigirka*, and *Kolima*. In the narrower belt of land which falls to the opposite seas, the chief rivers are the *Zambezi* in Africa; the *Euphrates*, *Indus*, *Ganges*, and *Brahmaputra*,

* See Philosophical Transactions, 1833. Rev. W. Whewell, M.A., F.R.S., &c., on "An approximation to a map of cotidal lines."

and the *Irrawadi* in southern Asia ; with the *Murray* river in Australia, flowing to the Indian Ocean. The *Mekong*, *Canton River*, *Yangtse*, *Hoang Ho*, and *Amur* in eastern Asia ; the *Columbia* and *Colorado* in North America, are the main streams which flow to the Pacific. The little streams which reach the Pacific from the steep western side of South America, contrasted with the mighty rivers which flow over its gentle slope to the Atlantic, give an extreme type of the two great ocean systems.

Areas of interior drainage are distributed over the whole of the continents, and are found centering in those drier regions, which, either from their remoteness from the oceans which supply moisture to the rain-bringing winds, or from their physical position—it may be screened from rain by an intervening chain of mountains which exhausts the rainfall on its sea-facing side—have not a sufficient deposition of moisture within them to form rivers large enough to resist the evaporation in the dry region through which they must pass to reach the sea. The terminal point of continental rivers is generally an inland sea or lake, the waters of which have become salt through the gradual accumulation of saline particles brought to it by its feeders. In these lakes a more or less perfect balance has been arrived at between evaporation and river supply, and the uniform extent of the lake is thus maintained. The continental basin of Asia, of which the *Caspian*, the *Sea of Aral*, *Lake Balkash*, and *Lake Lob* are the central points, is the largest of all, and the *Volga* is the greatest river of any interior basin. In Africa there are two extensive continental basins, of which *Lake Chad*, in the north, and *Lake Ngami*, in the south, are the centres. The whole of Central Australia may be considered as a continental basin, having its representative salt lakes in *Lakes Eyre* and *Torrens*. A basin of interior drainage surrounds the great *Salt Lake* of North America. Mexico and the Argentine Republic of South America have also continental basins.

The areas of the river systems belonging to each of the oceans, and of the continental basins, have been estimated as follows* :—

	Square Miles.
Area draining to the Atlantic Ocean	19,050,000
„ „ Arctic Ocean	7,500,000
Drainage area of the gentler slopes of the continents	26,550,000
Area draining to the Pacific Ocean	8,660,000
„ „ Indian Ocean	6,300,000
Drainage area of the steeper slopes of the continents	14,960,000
United areas of continental drainage	10,673,000

The size of a river may be variously estimated by the measurement of its length from source to mouth along its windings, by the extent of its collecting basin, by the volume of water which it discharges to the sea at its mouth, or by the extent of navigable channel which it affords. But there can be no one satisfactory standard of comparison, since two rivers which have the same length may differ very much in the extent of their tributaries ; or two rivers which have basins of the same area may be

* From Keith Johnston's Handbook of Physical Geography.

situated, the one in a dry, the other in a moist region, in which case the volumes of water collected by each will differ exceedingly. In every respect, however, the *Amazon*, which has been called the "Mediterranean of South America," is the greatest river of the globe, affording a water-way from the Atlantic to the very base of the Andes. The great rivers which come next to it, in the order of the extent of their basins, are given in the following table:—

	Sq. Miles.		Sq. Miles.
Amazon	2,200,000	St. Lawrence	1,099,000
Obi	1,295,000	Yenisei	896,000
Parana and Paraguay	1,250,000	Amur	870,000
Mississippi	1,235,000	Yangtse	845,000
Nile	1,100,000 (?)		

The chief fresh water lakes of the globe, arranged in the order of the systems to which they belong, may be compared as follows, in extent and elevation above the sea:—

1.—ARCTIC SYSTEM.

	Area in sq. miles.	Height in feet.
Great Slave Lake, Mackenzie R.	10,800	400 (?)
Great Bear Lake	7,000	250
L. Zaizan, in the basin of the Obi	1,500	1,918
L. Baikal, Yenisei basin	13,300	1,280

2.—ATLANTIC SYSTEM.

L. Winnipeg, Nelson R.	9,000	628
L. Superior, St Lawrence R.	32,000	627
L. Michigan, "	22,400	578
L. Huron "	22,000	578
L. Erie "	11,000	565
L. Ontario "	6,300	231
L. Nicaragua, S. Juan R., Cent. Amer.	7,170	130
L. Maracaybo, Venezuela	7,800	near sea level.
L. Tsana or Dembea, Blue Nile, Abyssinia	1,360	6,270
Victoria Nyanza, Upper Nile	30,000 (?)	3,740 (?)
Albert Nyanza "	25,000 (?)	2,500
L. Bangweolo, Upper Congo R.	6,000 (?)	4,000
L. of Geneva, R. Rhine	221	1,230
L. Constance, R. Rhine	183	1,290
L. Onega, R. Neva	3,400	237
L. Ladoga "	7,150	49
L. Wetter, R. Motala, Sweden	714	290
L. Wener, R. Gotha, "	2,015	143
L. Neagh, R. Bann, the largest in the British Isles	154	48

3.—PACIFIC OCEAN.

L. Lanquihue, S. Chili	320	172
Tulare Lake, R. Sacramento, California	550	398
L. Hinka, R. Amur	1,550	2,000 (?)
L. Toongting, R. Yangtse, China	2,340	200 (?)
Thale Sab, R. Mekong, Cambodia	1,100	50 (?)
L. Nyassa, Zambezi, N. Africa	8,500	1,522

The lakes of continental basins are almost invariably salt or brackish where they have no outlet to another lake in the same basin; though there are some exceptions to this rule. Among the chief are,—

The Caspian Sea	178,866	84 (below sea)
Sea of Aral	26,940	26 (above)

	Area in sq. miles.	Height in feet.
L. Balkash, Central Asia	11,500	700 (?)
Issyk-Kul „	2,466	4,480
L. Sistan, Persia	900	1,545
L. Van, Armenia	1,413	5,465
L. Urumiah „	1,730	4,000
Dead Sea, Palestine	495	1,298 (below)
L. Chad, Soudan (fresh)	14,500	830 (?)
L. Tanganyika, Cent. Africa (fresh)	10,400	2,880
L. Ngami, S. Africa	300	2,800
Salt Lake of Utah, Wn. N. America	1,630	4,210
L. Titicaca, Bolivian plateau	3,200	12,846
L. Eyre, S. Australia	4,000 (?)	70

PLATES XIII, XIV.—Hydrographic Map of the British Isles.

The inland waters of the British Isles distribute themselves naturally, according to the cardinal points, into Northern, Eastern, Southern, and Western systems. The boundaries of these systems, the great water-partings of the islands, are shown on the map by strong red lines, while the minor divisions of these systems into river-basins are indicated by lighter lines of the same colour. The lengths and drainage areas of all rivers whose basin exceeds 400 square miles are given on the map.*

The elongated form of the island of Great Britain, in the direction from north to south, causes the line dividing the eastward from the westward flowing waters to be the most important in the island; and since the more elevated parts of the land are in the west, this line, following the highest ground, lies uniformly nearer to the west coast than the east. In Ireland the main water-parting is that which runs from north-west to south-west in a curve approaching the east coast in its centre, showing that, contrary to the case of Great Britain, the greater drainage of the land is to the westward. More than one-half of the area of Britain is drained to the eastward, and nearly one-half of the area of Ireland to the westward.

The following table gives the areas of the different systems of the British Isles in English square miles:—

	Northern.	Southern.	Western.	Eastern.
England,	...	7,306	21,255	29,759
Scotland (Mainland), .	1,120	2,700	7,637	15,356
Great Britain, . .	1,120	7,306	31,592	45,115
Ireland,	4,800	6,800	15,750	5,150
British Isles, . . .	5,920	14,106	47,342	50,265

From this table it will be seen that the Southern system of England is very small in comparison with its Western and Eastern systems, being only one-third of the former

* The lengths of the rivers and areas of the river-basins of England given have been taken from the "Plan of the Catchment Basins of the Rivers of England and Wales," published at the Ordnance Survey Office in 1861. Those for Scotland and Ireland are the results of a careful series of original measurements from large-scale maps.

and one-fourth of the latter in area ; and that the smallest system of all in the islands is that of the north of Scotland, whose extent is only a fortieth part of that of the greatest, the Eastern system of Britain.

If Scotland be considered by itself, separately from England, it may be said to have a southern drainage system of 2700 square miles in extent to the Irish Sea ; but if taken in connection with England, as forming with it the island of Great Britain, this area must be looked upon as a part of the Western drainage system of the island. The Northern, Southern, and Eastern systems of Ireland are of nearly equal size, and the three taken together form an area not much greater than that of the Western system.

The Eastern system of Great Britain, included within the water-parting line which runs from Duncansby Head, on the north-east, to near Dover, on the south-east, and occupying more than one-half of the area of the island, embraces most of the larger rivers of Britain, and discharges its waters to the North Sea. Eleven of the river-basins of this system have an area of upwards of 1000 square miles, and are those of the Spey, Tay, and Tweed, in North Britain ; and of the Tyne, Ouse, Humber, Trent, Witham, Nen, Great Ouse, and Thames in South Britain—the last-named, the Thames, being the greatest river of the British Isles, and draining an area of 5162 square miles.

The Western system of Great Britain, stretching from Cape Wrath to Land's End, and draining to the Atlantic and the Irish Sea, has but four river-basins of greater extent than 1000 square miles, these being the Clyde in Scotland, and the Mersey, Wye, and Severn in England. The northern part of this system has no rivers, properly so called, but is drained by numerous streams, which rise with rain and fall with dry weather ; but in the south of it, the Severn, with its basin of 4437 square miles, is the third river of the British Isles.

The Southern system of Britain is a narrow strip of land extending between Dover and Land's End, draining to the English Channel, and having no great rivers—the largest basin, that of the Avon, being only 666 square miles in extent.

The Northern system in Britain is one of streams alone ; the largest, the Naver, having a basin of only 200 square miles.

The Southern, Western, and Northern systems of Ireland send their waters to the Atlantic. The Eastern system drains to the Irish Sea.

The Western system of Ireland, like the Eastern of Britain, embraces about one-half of the area of the island. The water-parting bounding it runs from Bloody Foreland, in the north-west, to Mizen Head, in the south-west, including three river-basins of upwards of 1000 square miles in area—those of the Erne, Galway, and Shannon ; the last is the largest river of Ireland, and the second of the British Isles, draining an area of 4590 square miles. The water-parting of the Southern system of Irish rivers runs inland to near the centre of Ireland, round the head-waters of the Barrow, its largest river, and meets the coast at Carnsore Point on the east, and Mizen Head on the west. Besides the Barrow, the Blackwater is the only other river of this system with a basin of more than 1000 square miles.

The Eastern system of Ireland extends in a narrow strip from Fair Head, on the

north, to Carnsore Point, in the south. Its only river-basin of above 1000 square miles in area is that of the Boyne.

Almost the entire area of the Northern system of Ireland which lies between Fair Head and Bloody Foreland is occupied by the rivers Bann and Foyle, each of which drains an area of upwards of 1000 square miles.

In the order of the extent of their basins, the rivers which drain an area of upwards of 1000 square miles stand as follows:—

	Sq. miles.		Sq. miles.
1. Thames,	5162	13. Wye,	1655
2. Shannon,	4590	14. Blackwater,	1300
3. Severn,	4437	15. Spey,	1245
4. Ouse,	4207	16. Humber,	1229
5. Trent,	3972	17. Galway,	1172
6. Barrow,	3517	18. Clyde,	1145
7. Great Ouse,	2894	19. Foyle,	1090
8. Bann,	2265	20. Nen,	1055
9. Tay,	2090	21. Tyne,	1053
10. Tweed,	1990	22. Witham,	1052
11. Mersey,	1706	23. Boyne,	1046
12. Erne,	1660		

The contour lines on the map are of 250, 500, 750, and 1000 feet, and have been worked out chiefly from the books of the Ordnance Spirit-Levelling Survey. Though they can only be said to be an approximation to the truth, yet they give a much more accurate representation of the elevation of the land than can be obtained by any system of hill-shading.

From these lines an estimate may be made of the elevation of the source and the average rapidity of fall of any river, by observing its passage from one height to another. If the contours approach closely to one another in a river-course, its fall is most probably rapid, and the river unfit for purposes of navigation; but if, on the contrary, the lines be far apart, and at equal distances from each other, we may predicate that the river has a gentle descent to the sea.

The blue tints representing the rainfall have been obtained by taking the mean of the rainfall, as given in Mr Symons's reports* for the five years from 1863 to 1867 inclusive, of nearly 500 stations in the British Isles. The figures thus found were set down on large-scale maps, and lines were drawn through the places which have the same average rainfall, whilst keeping in view the general fact that a greater elevation gives a greater rainfall.

From these tints it is evident that the western parts of the islands receive the greatest amount of rain. This may be due to two causes—first, to the prevalence of rain bringing south-west winds; and second, and perhaps chiefly to this, that the western parts of the islands are generally of greater elevation than the eastern.

The part of the British Islands which shows the least rainfall is that strip of the east coast which lies between the mouth of the Humber and the Wash, the rainfall there being under 20 inches. A small portion of the north coast of Elginshire has also, perhaps, this minimum of rainfall.

* "British Rainfall." Compiled by G. J. Symons, F.M.S.

Almost the whole of eastern England has a rainfall of from 20 to 25 inches—the Downs in the south and the Wealds in the north being the exceptions; but only some small patches of the east coast of Scotland have this minimum rainfall, and the fall in all parts of Ireland seems to be greater than this.

The line of 30 inches of rainfall runs as far back as the Welsh boundary in Mid-England, but is carried eastwards again by the Pennine chain and the Cheviots, and includes only the east coast of Scotland, though brought considerably inland by the valley of the Tweed, the low land between the rivers Forth and Clyde, and Strathmore of the Tay Basin.

The lowest rainfall in Ireland, as far as has yet been ascertained, is from 25 to 30 inches on the lower parts of the south-east and east coasts. The central plain of Ireland lies entirely within the lines of 30 to 35 inches of fall. The lines of 35 to 45 inches and above are confined to the higher parts of west and south-west England, and the Pennine chain in the north, but include the great mass of Scotland and the whole of Ireland, excepting the central plain and the east coast. The observing stations on the west coasts and in the hilly regions are not sufficiently numerous to enable the lines of greater rainfall to be drawn on the Map; but it is most probable that these lines would be found to follow the same law which holds good in the other parts of the islands, of increase with greater elevation and proximity to the western coasts.

The maximum observed rainfall seems to be that noted at Seathwaite, in the Cumberland mountains, which, in the mean of the five years from 1863 to 1867, has a fall of 140 inches; but many of the higher parts of the west of England, Scotland, and Ireland have doubtless a rainfall of more than 100 inches.

PLATE XV.—Rain and Snow.

Rain is distributed very unequally over the globe. In general it is most abundant in those regions where evaporation is carried on most rapidly; but there are striking exceptions to this rule, for in many places, even near the tropics, it seldom or never rains. These almost rainless districts comprise vast regions, nearly devoid of vegetation. On the contrary, there are regions where rain falls almost incessantly, and where, consequently, vegetation is rank and abundant. The zone of greatest precipitation is situated on the north of the Equator, and corresponds with the zone of "the variables" in Plate XVII. Within this belt, on the west coast of Africa, the average annual quantity is 189 inches. Rain is in general most abundant near the Equator, and the quantity decreases, irregularly, with the decrease of heat, as we approach the polar regions. Within the tropics the rains follow the apparent movement of the sun; thus countries lying towards the tropic of Cancer have their rainy seasons in the summer of the northern hemisphere, those lying near the southern tropic, during our winter; and since the sun passes the Equator twice during the year, there is a double rainy season along the line. The average annual fall within the tropics is 95 inches, and within the temperate zone, 34 inches. This is shown graphically in the diagram ("Increase of Rain with Heat") at the foot of the Plate. The diagram of "Decrease of Rain with Distance from Coasts," illustrates a general law occasioned by the greater

amount of vapour that arises from the sea than from the land. In the temperate zone of both hemispheres, the western coasts are proportionally more moist than the eastern, because they are exposed to the prevailing westerly currents of air, which, passing over the ocean, are highly impregnated with moisture. Within the tropics, the eastern coasts, especially in America, are more moist than the western, from their exposure to the trade winds. In certain districts more rain falls in one season than another : these are distinguished on the Map as the summer, winter, and autumn rains. The almost rainless areas of the globe are due to the drying of the winds which reach these regions, either from their passing from a colder to a warmer zone, as in the case of the Sahara, or from their having expended their moisture on the margins of the continents, as in Central Asia and the great basin of North America.

Snow never falls within or near the tropical regions at the level of the sea. The red lines on the Map show that the deposition of moisture in this form is, in the northern hemisphere, limited to the regions north of India, the Mediterranean, and the Gulf of Mexico ; while in the southern hemisphere it does not approach nearer the Equator than the south of Australia, the Cape Colony, and Patagonia. But in tropical and sub-tropical regions snow falls on mountains where, owing to elevation above the sea, the temperature is reduced to the freezing point ; as in the Himalayas in Asia, Mounts Kenia and Kilimandjaro in Eastern Africa, and the Andes of South America. The fall of snow increases with the decrease of temperature, in proceeding from the equatorial towards the polar regions. In Europe this decrease occurs in the following order :—Rome has $1\frac{1}{2}$ snowy days in each winter, Venice $5\frac{1}{2}$, Milan 10, Paris 12, Carlsruhe 26, Copenhagen 30, and St Petersburg 171. The actual elevation of the snow line—the line of perennial congelation—or the lowest point on a mountain at which snow is never entirely melted, in the different zones, is shown in the diagram on the left of the Map, and on an enlarged scale in Plates XVIII. and XIX.

PLATE XVI.—Climatology*—Isothermal Lines.

The distribution of the sun's heat over the surface of the globe is here rendered perceptible to the eye by lines drawn through all places having the same mean annual temperature. These lines are called *isothermals*, and their deviation from parallelism with the Equator shows that latitude, or distance from the equator (or from the more direct power of the sun) alone, gives little indication of the temperature of any particular place, since many causes unite to disturb their arrangement.

The mean annual temperature of the more important places on the globe is stated in figures on the larger Map. The Map of Central Europe shows by red and blue figures the summer and winter temperatures of the principal meteorological stations, while that of the British Islands presents the three elements of summer, winter, and mean annual temperatures. If these places were connected by lines, the first would be called *isotheral*, the second *isochimnal*, and the third *isothermal* lines. July is, on an average, the hottest, and January the coldest, month of the year in the northern

* See Buchan's "Handy Book of Meteorology," 2d Edition, Edinburgh, 1868, where the whole subject of Climatology is treated of in an interesting and lucid manner.

hemisphere, these conditions being reversed in the southern. The mean annual temperature occurs in April and October in the north temperate zone. On each side of the meridian of 30° west, the extreme temperature of summer and winter (July and January), for the latitude, is stated in red figures for summer, and blue for winter. A remarkable example of the disturbance of the latitudinal arrangement of the isotherms occurs on the west coast of Europe, where the warm water of the Gulf Stream has the effect of carrying the lines many degrees north of their normal position, whilst they are pushed southward by the cold currents of the American coast, so that London enjoys an average temperature of 51° whilst places on the American coast in the same latitude have that of the freezing point. A climate is called an *insular* or *sea* climate where the difference of mean temperature is very small, or where the winter is milder and the summer cooler than the average; and a *continental* climate where the difference of mean temperature is very great, or where the winter is very cold and the summer very hot. Thus the differences between the summer and winter temperature of a country increase with distance from the sea. Europe has a true insular climate—a mild winter and a cool summer. Northern and Central Asia have a true continental climate—a cold winter and a hot summer; while North America has more of a continental climate in winter, and a sea climate in summer. The hottest locality on the globe is in northern Central Africa (shaded red on the Map), where the temperature of July is 90° Fahr.; and the coldest in Siberia (shaded blue on the Map), where the temperature of January is 40° below the freezing point of Fahr. scale. The cold region of Siberia has no corresponding region of equal cold in America.

If the globe be divided at the meridian of 20° W., we find that the eastern portion, which has the largest mass of land, is colder than the western, and that the difference diminishes as we approach the Equator. The temperature of the whole globe is believed to increase 8° Fahr. from January to July; a mean between these months gives, as the mean temperature of the globe, 58° Fahr. The mean temperature for the northern hemisphere is 60° Fahr., and the mean for the southern hemisphere 56° Fahr. The great quantity of rain which falls in the northern hemisphere is probably one cause of its higher temperature, while, in the southern hemisphere, the influence of the sun's rays is expended to a great degree in the melting of masses of ice, or in evaporation from the surface of the ocean.

PLATE XVII., Winds and Storms,

presents a graphic representation of the principal phenomena of the winds over the globe. Near the centre is the great equatorial belt of calms and variable winds and storms, corresponding to the zone of constant rain in Plate XV., and on the north and south of the trade-wind region, the narrower belts of the calms of Cancer and Capricorn. The region of the *Trade* or passage winds, called *Vents alisés* by the French, extends to about 30° on each side of the Equator, and beyond these are the regions of the south-westerly and north-westerly currents of air. The Indian Ocean is the chief region of the *Monsoon winds* (from the Arabic "Mausim," an epoch or season), included within which is the district of the *Typhoons*, or storms peculiar to the China Sea. But monsoons, or monsoon-like winds, prevail to a limited extent in various

parts of the globe, as on the shores of Brazil, Chile, Peru, Mexico, and the west and north coasts of Africa, indicated by the red colour on the chart. The Red Sea and the Persian Gulf, not being affected by monsoon winds, are left white.

The diagrams at the foot of the Plate are devoted to an explanation of the *Hurricanes* of the West Indies and the Indian Ocean, and of the varying limits of the trade-winds in the Atlantic, according to the different seasons of the year. The hurricanes of the Indian Ocean and of the West Indies, and the typhoons of the China Sea, appear to be subjected to peculiar and fixed laws for each hemisphere, both as regards their movements of translation and their rotatory movements. These tempests originate between the Equator and the tropics during winter, when the regularity of the trade-winds is interrupted, or during the change of the monsoons. They obey a double movement—the one of translation or progress, and the other of a gyratory or rotatory kind. On the north of the Equator the rotatory movement is from *right* to *left*, while on the south of the Equator it is reversed, or passes from *left* to *right*, as shown on the Plate. The movement of translation takes the form of a parabolic curve, the summit of which is towards the west. In the northern hemisphere this curve forms a tangent to the meridian about lat. 30° , and in the southern hemisphere about lat. 26° , corresponding with the northern and southern limits of the trade-winds. In the Indian Ocean, these storms uniformly come from the eastward, and travel to the westward and southward, as shown in the chart of the Rodriguez hurricane. The West Indian hurricanes commence near the Leeward Islands, travel to the W.N.W., and then round the shores of the Gulf of Mexico, following the course of the Gulf stream, and are lost in the Atlantic between the Bermudas and Halifax. The chart in the left corner of the Plate shows the dates of occurrence, and the courses, of some of the most important of these storms. The rate of progress of a hurricane varies in different parts of its course; that of the Rodriguez storm has been calculated at from 220 to 230 miles a-day at first, diminishing, as it approaches the tropics, to about 50 miles a-day.

The districts subject to local storms are marked in the Map as the *Sirocco* in the Levant, the *Simoom* in Arabia, the *Khamsin* in north-east and the *Harmattan* in south-west Africa, these being different names for the hot winds which blow out from the deserts of the Sahara at various seasons, the *Pamperos*, sudden gusts of S.W. wind in South America, the *Norther*s or cold blasts of N. wind in Mexico, and the *Hot Winds* in Australia.

PLATE XVIII.—Vegetable Life.

The plants of any particular region are the exponents of its climate: certain plants will grow spontaneously only within certain districts or zones, the boundaries of which are dependent on the amount of heat and moisture which such zones receive in the course of the year. Hence we find that trees, grain, and shrubs range themselves on the globe according to lines of equal mean summer and equal winter temperature: the lines of summer temperature, for example, indicate precisely the limits of the possible cultivation of annual plants. Nor is the knowledge of the

capabilities of a country for producing plants less important with reference to its population. Comparing Naples with Norway, for example, we find that the effect of climate is such as to render the harvest five times more productive in the former than in the latter place: while, in consequence, the population is twenty-five times more dense, in proportion to its area, in Naples than in Norway.

The object of this Map is, by dividing the whole earth according to its peculiar flora into certain climates of vegetation, to present at one view the distribution of the most useful and valuable wild and cultivated plants. Each of the plant-climates is characterised by certain trees, grains, and fruits, the number and variety of which increase as we approach the Equator, and decrease towards the polar regions, where their only representative are a few mosses and grasses. Rice, which supports the greatest number of the human family, is chiefly confined to the tropical regions of Asia and America. Wheat requires a mean annual temperature of 37° or 39° Fahr., and a mean summer temperature of 56° to 58° at least. Maize extends to lat. 40° N. and S. in America, and to lat. 50° or 52° N. in Europe; barley to lat. 70° N. in Norway; rye to lat. 67° , and oats to lat. 65° N. Since heat decreases as we ascend into the atmosphere, so mountains situated between the tropics, the summits of which rise above the snow-line, represent the vegetable zones of the whole earth, rising one above the other *vertically* in the same order as they observe in a *horizontal* direction from the Equator to the Poles, on plains. This is explained by the colours on the diagram at the bottom of the Plate, on which also is represented the actual ascertained limits of the snow-line in different latitudes. In the Himalaya this will be observed to be 19,000 feet on the north, while it is only 15,500 feet on the south side—an anomaly which has been considered due to the radiation of heat from the high land of Tibet, but which is more probably owing to the greater quantity of snow which falls on the southern than on the northern slopes of the mountains.

PLATE XIX.—Animal Life.

Animals, like plants, are adapted to special climatic conditions; like them, also, they are subjected to invariable laws. Each zone, or region of climate, is occupied by some species of animals peculiar to itself, beyond the limits of which they will not range if left to their natural freedom. A group of animals embracing all the species both terrestrial and aquatic, inhabiting any particular region, is called a *Fauna*, in the same way as the plants of any particular country constitute its *Flora*. In this Map the faunas of the two hemispheres are distributed in three principal divisions—namely, the Tropical, Temperate, and Arctic faunas, each of which is characterised by peculiar species of animals. These, like plants, attain their highest development within the torrid zone, which is distinguished not only by the great variety, but by the size strength, and beauty of its animal formations. The mean density of species inhabiting, the different zones of the earth strikingly proves this. The proportion is estimated thus: in the tropical region, 26 species; in the north-temperate, 9; and in the arctic, only 7 of the whole. In proportion as they recede from the Equator, so the covering of animals loses its brilliancy of colour, until, in the polar regions, all animals, whether

marine or terrestrial, assume a nearly uniform and sombre hue. From a comparison of this with the preceding Map, it will appear that the nature of their food has an important bearing on the distribution and the grouping of animals, those of the herbivorous kind being more or less limited to particular vegetable zones, while, since the food of carnivorous animals is everywhere present, their range is, in this respect, much less confined, the absence of animal life itself being the only restriction.

The diagram at the foot of the Plate represents the distribution of animals in a vertical direction, showing, as in the case of plants, that those nearest the level of the sea correspond with those of the torrid zone; while, on approaching the snow-line near the tops of mountains, the animals are similar to those of the arctic zone.

PLATE XX.—Races of Man.

The geographical distribution of man is different from that of all other organic beings: his constitution renders him more cosmopolite, while, from the superior structure of his physical frame, as well as his mental endowments, he is less subject to the influence of external circumstances than any of the inferior animals: hence he is found, although not in equally favourable circumstances, in every locality over the globe, under every climate, and at every degree of altitude to which organic life extends. All the various races of man existing on the globe have originated from one species, but distinct races have been known to exist since the earliest dawn of history and tradition. The principal Map exhibits the distribution of the leading races of man, as existing at the present time, the localities of which are explained by the reference figures and colours.

The Ethnographic Map of Europe presents a more minute sub-division of these races in this quarter of the world. It distinguishes by colours three great varieties of the Caucasian division—the Teutonic, the Celtic, and the Slavonian. The sub-varieties have different shades of these fundamental colours; and wherever there has been a crossing of these varieties, it is indicated by a mixed tint. The diagram in the right corner of the Plate explains, by means of the scale of feet, the height above the level of the sea of inhabited places in different countries. Coloured lines and a scale placed near the meridian of 20° W. on the upper Map, indicate the prevailing kinds of food used by the inhabitants of the different zones. Thus within the tropics the principal food of the human family is derived from the vegetable kingdom; beyond the tropics, north and south, a mixed animal and vegetable diet is used; while north of the Arctic circle the food of man is derived entirely from the animal kingdom. A faint line passing through Greenland and the north of Asia points out the northern limit of permanent habitation; the corresponding line in the southern hemisphere passes south of Cape Horn and Tasmania.

ELEMENTARY WORKS

PUBLISHED BY

WILLIAM BLACKWOOD & SONS.

NATURAL SCIENCE CLASS-BOOKS.

A MANUAL OF ZOOLOGY, for the Use of Students. With a General Introduction on the Principles of Zoology. By HENRY ALLEYNE NICHOLSON, M.D., D.Sc., M.A., F.R.S.E., &c., Professor of Natural History in the University of Toronto. Second Edition. Crown 8vo, pp. 674, with 243 Engravings on Wood, price 12s. 6d.

TEXT-BOOK OF ZOOLOGY, for the Use of Schools. By the same Author. Crown 8vo, with 153 Engravings on Wood, price 6s.

INTRODUCTORY TEXT-BOOK OF ZOOLOGY, for the Use of Schools. By the same. A New Edition. 127 Engravings, price 2s. 6d.

INTRODUCTORY TEXT-BOOK OF GEOLOGY. By DAVID PAGE, LL.D., F.R.S.E., &c. With Engravings on Wood and Glossarial Index. Ninth Edition, price 2s.

ADVANCED TEXT-BOOK OF GEOLOGY, Descriptive and Industrial. By the same. With Engravings and Glossary of Scientific Terms. Fifth Edition, revised and enlarged, price 7s. 6d.

GEOLOGY FOR GENERAL READERS. A Series of Popular Sketches in Geology and Palaeontology. By the same. Third Edition, enlarged, price 6s.

HANDBOOK OF GEOLOGICAL TERMS, GEOLOGY, AND PHYSICAL GEOGRAPHY. By the same. Second Edition, enlarged, price 7s. 6d.

INTRODUCTORY TEXT-BOOK OF PHYSICAL GEOGRAPHY. With Sketch-Maps and Illustrations. By the same. Fifth Edition, price 2s.

ADVANCED TEXT-BOOK OF PHYSICAL GEOGRAPHY. By the same. With Engravings, price 5s.

HISTORICAL CLASS-BOOKS.

EPTOME OF ALISON'S HISTORY OF EUROPE, for the Use of Schools and Young Persons. Post 8vo, pp. 604, price 7s. 6d., bound in leather. Sixteenth Edition.

HISTORY OF FRANCE FROM THE EARLIEST TIMES TO 1848. By the Rev. JAMES WHITE, Author of 'The Eighteen Christian Centuries.' Post 8vo, with Index, price 6s.

THE EIGHTEEN CHRISTIAN CENTURIES. By the Rev. JAMES WHITE, Author of 'The History of France.' School Edition, post 8vo, with Index, price 6s.

ANCIENT CLASSICS FOR ENGLISH READERS. Edited by the Rev. W. LUCAS COLLINS, M.A. Price 2s. 6d. bound in cloth. A Volume is published Quarterly.

THE SCHOOL ETYMOLOGICAL DICTIONARY AND WORD-BOOK. Combining the advantages of an ordinary Pronouncing School Dictionary and an Etymological Spelling-book. By the Rev. JAMES STORMONTH. Fcap. 8vo, pp. 254, 2s.

GEOGRAPHICAL CLASS-BOOKS.

MANUAL OF MODERN GEOGRAPHY, Mathematical, Physical, and Political, on a New Plan; embracing a complete Development of the River Systems of the Globe. By the Rev. ALEXANDER MACKAY, LL.D., F.R.G.S. A New and Improved Edition. Crown 8vo, pp. 676, 7s. 6d.

ELEMENTS OF MODERN GEOGRAPHY. By the same. Twelfth Edition. Crown 8vo, pp. 304, price 3s.

OUTLINES OF MODERN GEOGRAPHY. A Book for Beginners. By the same. Fourteenth Edition. 18mo, pp. 112, price 1s.

SCHOOL ATLASES BY A. KEITH JOHNSTON, LL.D., F.R.S.E., Author of the Royal and Physical Atlases.

ATLAS OF GENERAL AND DESCRIPTIVE GEOGRAPHY. A New and Enlarged Edition. 26 Maps, with Index. Half-bound, 12s. 6d.

ATLAS OF PHYSICAL GEOGRAPHY. A New Edition, revised and enlarged. Half-bound, 12s. 6d.

ATLAS OF CLASSICAL GEOGRAPHY. A New and Enlarged Edition. With Index. Half-bound, 12s. 6d.

ATLAS OF ASTRONOMY. A New and Enlarged Edition, 12s. 6d.

ELEMENTARY ATLAS OF GENERAL AND DESCRIPTIVE GEOGRAPHY, for Junior Classes. With Index. Half-bound, 6s.

THE THEORY OF ARITHMETIC. By DAVID MUNN, F.R.S.E., Mathematical Master, Royal High School, Edinburgh. Crown 8vo, price 5s.

A TREASURY OF THE ENGLISH AND GERMAN LANGUAGES. Compiled from the best Authors and Lexicographers. By JOSEPH CAUVIN, LL.D., Ph.D. of the University of Göttingen. Crown 8vo, price 7s. 6d.

HANDBOOK OF PHYSICS. By WILLIAM ROSSITER, F.R.A.S., &c., Author of 'Elements of Plane and Spherical Trigonometry,' &c. Crown 8vo, with Diagrams, 5s.

ENGLISH PROSE COMPOSITION: A Practical Manual for Use in Schools. By JAMES CURRIE, M.A., Principal of the Church of Scotland Training College, Edinburgh. Eighth Edition, 1s. 6d.

PROGRESSIVE AND CLASSIFIED SPELLING BOOK. By HANNAH R. LOCKWOOD, Authoress of 'Little Mary's Mythology.' Fcap. 8vo, 2s.

LATELY PUBLISHED,

NEW ATLAS FOR PUPIL TEACHERS

THE HANDY ROYAL ATLAS. 45 Maps clearly printed and carefully coloured, with GENERAL INDEX by A. KEITH JOHNSTON, F.R.S.E., F.R.G.S., &c. Imp. 4to, £2, 12s. 6d., half-bound morocco.

"The 'Handy Atlas' is thoroughly deserving of its name. Not only does it contain the latest information, but its size and arrangement render it perfect as a book of reference."—*Standard*.

A GEOLOGICAL MAP OF EUROPE, exhibiting the different Systems of Rocks according to the latest researches, and from Inedited materials. By SIR R. I. MURCHISON, D.C.L., F.R.S., &c., Director-General of the Geological Survey of Great Britain and Ireland; and JAMES NICOL, F.R.S.E., F.G.S., Professor of Natural History in the University of Aberdeen. Constructed by ALEX. KEITH JOHNSTON, F.R.S.E., &c. Scale, $\frac{1}{4,800,000}$ of Nature, 76 miles to an inch. Four Sheets Imperial, beautifully printed in Colours. Size, 4 feet 2 inches by 3 feet 5 inches. In Sheets, £3, 3s.; in a Cloth Case, 4to, £3, 10s.

ETYMOLOGICAL AND PRONOUNCING DICTIONARY OF THE ENGLISH LANGUAGE. Including a very Copious Selection of Scientific Terms. For Use in Schools and Colleges, and as a Book of General Reference. By the Rev. JAMES STORMONTH. The Pronunciation carefully Revised by the Rev. P. H. PHELP, M.A., Cantab. Crown 8vo. pp. 755. 7s. 6d.

HANDY BOOK OF METEOROLOGY. By ALEXANDER BUCHAN, M.A.; F.R.S.E., Secretary of the Scottish Meteorological Society. A New and Enlarged Edition, being the Third. *[In the Press.]*

"A very handy book this, for in its small compass Mr Buchan has stored more and later information than exists in any volume with which we are acquainted."—*Symon's Meteorological Magazine*.

"Clear, concise, and easy of reference."—*The Field*.

"We do not know a better work on Meteorology."—*Gardener's Chronicle*.

THE ORIGIN OF THE SEASONS, Considered from a Geological Point of View, showing the Remarkable Disparities that exist between the Physical Geography and Natural Phenomena of the North and South Hemispheres. By SAMUEL MOSSMAN, Author of 'China, Its History and Institutions,' 'Our Australian Colonies,' &c. One Volume, with Maps and Diagrams, 10s. 6d.

THE PAST AND PRESENT LIFE OF THE GLOBE. By DAVID PAGE, LL.D., F.R.S.E., F.G.S. With numerous Illustrations. Crown 8vo, 6s.

COMPARATIVE GEOGRAPHY. By CARL RITTER, Professor of Geography in the University of Berlin. Translated by W. L. GAGE. Fcap. 8vo, price 3s. 6d.

TEXT BOOK OF ZOOLOGY, for the Use of Schools. By HENRY ALLEYNE NICHOLSON, M.D., D.Sc., M.A., Ph.D., Göt., F.R.S.E., F.G.S., &c., Professor of Natural History in the University of Toronto. Crown 8vo, with Engravings on Wood, price 6s.

"The author's Text-book, regarded from this point of view, is a decided success; it is just what was wanted. The subject has been treated in a scientific spirit, but at the same time so clearly and well, as to be quite within the comprehension of any young student who will bring ordinary attention to his task."—*Lancet*.

"If the book is solely intended for the use of schools, it is just such a book as a boy may find time to thoroughly master; a stepping-stone, as it were, to larger treatises on the same subject, and in this way will prove both acceptable and useful."—*Examiner*.

"This is an excellent treatise for its purpose. Dr Nicholson's style is singularly intelligent; he conveys much information in a small space, and is especially clear in his explanations of the various zoological classes and their characteristics. The introductory chapter is a model of lucid writing on a difficult point."—*Nonconformist*.

"The best Advanced Text-book of Zoology that has yet appeared."—*The Student*.

BY THE SAME AUTHOR,

INTRODUCTORY TEXT-BOOK OF ZOOLOGY, for the Use of Junior Classes. A New Edition. With Illustrations. Price 2s. 6d.

BY THE SAME AUTHOR,


A MANUAL OF ZOOLOGY, for the Use of Students. With a General Introduction on the Principles of Zoology. Second Edition. Crown 8vo, pp. 674, with 243 Engravings on Wood, price 12s. 6d.

"A work which, in point of accuracy, far surpasses most of the Manuals which of late years have issued from the press."—*Popular Science Review*.

"It is, indeed, worthy of very high praise. In completeness and accuracy, as to the parts of the science of zoology of which it treats—the structure and classification of animals—it excels every work of the same kind with which we are acquainted."—*Courant*.

WILLIAM BLACKWOOD & SONS, EDINBURGH AND LONDON.

ILLUSTRATIONS OF THE ACTION OF RAIN AND STREAMS



2



3



7



4



8



5

Diagram section of a Waterfall

a Soft shales which decompose more rapidly than the rock above
b Hard limestone which projects beyond the soft rock below and gives rise to the perpendicular fall.




6

Serpentine Course of a River on flat ground


The river cuts away its banks most at the points marked *a* & *b*; sediment is heaped up at *c* & *d*, the connecting ground between two loops of the river. *e* is sometimes cut through giving rise to islands *f* & *g*, which remain as such until the old river course is filled up.



9



9a




9b

ILLUSTRATIONS OF THE ACTION OF ICE AND SNOW



ILLUSTRATIONS OF SEA ACTION.




SECTIONS ACROSS THE BASIN OF THE ATLANTIC OCEAN.



ILLUSTRATIONS OF VOLCANIC ACTION AND MOVEMENTS OF THE EARTH'S CRUST.




2 A STAGES IN THE GROWTH OF A CORAL-ISLAND



5 PROFILE OF A HILL-SLOPE SHOWING RAISED SEA-BEACHES






Engraved & Printed in Colours

William Blackwood & Sons, Edinburgh & London.

By W. A. K. Johnston, F.R.S.E.



TOPOGRAPHY




Drawn & Printed in 1870

William Blackwood





TOPOGRAPHY




Engraved & Printed in Colours

William Black




This is a detailed historical map of the Cape Colony and surrounding regions. The map is oriented with North at the top. It shows the Atlantic Ocean to the west, the Gulf of Guinea to the south, and parts of North and South Africa. Major cities and towns are labeled, including Cape Town, Stellenbosch, and Swellendam. The map also shows the Orange River and the Drakensberg mountains. A large inset map at the bottom shows the Cape Colony in more detail, with labels for 'ENLARGED MAP OF THE CAPE COLONY' and 'CAPE OF GOOD HOPE 1000'. The map is oriented with North at the top and includes latitude and longitude markings.




TOPOGRAPHY






This is a detailed topographical map of Central America and northern South America. The map is oriented with North at the top. The title "CENTRAL AMERICA" is prominently displayed in the center. The map shows the Isthmus of Panama, the Andes mountains, and various river systems. Key locations like Panama, Bogotá, and Caracas are marked. The map is oriented with North at the top, and the title "CENTRAL AMERICA" is prominently displayed in the center.





HYDROLOGY



Engraved & Printed in Colours

William Blackwood



WORLD.

aries, & extent of
VER BASINS
ifferent
S. LAKES, &c.
tribute their waters.
TON, F. R. S. E.

flow to the


Pacific Ocean. ☐
 Arctic Ocean. ☐

immediately reach the Sea.


Wilkes Land

On a scale corresponding with that of the Map.


E u r o p e A f r i c a & A s i a



by W. & A. K. Johnston, Edin.



HYDROGRAPHY



HYDROGRAPHIC MAP OF THE BRITISH ISLES

Showing the RIVER BASINS, the RAINFALL, and
the ELEVATION OF THE LAND in CONTOURS.

BY RICHARD JOHNSON, F.R.S.

Scale of English Miles
0 10 20 30 40 50 60 70

Explanation


The Boundaries and Names of the River Basins are in Red
The sloping figures after the name denote the length of the river and
the upright figures the Area of its Basin in sq. m. thus
The Rainfall is expressed in Inches according to this scale


The approximate contours are of
250 Feet thus
500
750
1000




Longitude West from Greenwich

William Blackie








T O R U N-P E R I O D I C A L R A I N S
 in the Year. & rain falls as often during night as day.


**DISTRIBUTION OF
 SNOW
 ON THE GLOBE.**
 Cause of Moisture
 REARDS THE POLAR REGIONS
 CAL AND SEASONAL RAINS
 WHEN NO RAIN FALLS
 IN THE POLAR REGIONS
 Water is the amount of Rain
 at the earth's surface, for a
 uniform fall of different Places
 the comparative amount of Rain
 falling in the present 1874.

Increase of Rain with heat.



The comparative amount of Rain
 which falls in the different Zones.


Decrease of Rain with distance from Coasts.



The decrease of Rain from the Coasts
 to the interior of Continents


1

METEOROLOGY



Engraved & Printed in Colours

William Blackwo.



CLIMATOLOGICAL CHART.

Showing by Lines & Figures
THE MEAN ANNUAL TEMPERATURE
of the principal places on the Globe.


BY A. K. JOHNSTON, F.R.S.E.


Explanation

The Climate Lines or Isotherms, connect places having the same mean annual temperature, those coloured Red, being above & those coloured Blue, below the Freezing point of Fahrenheit's scale.

The Red Figures denote Summer (July), the Blue, Winter (January), and the Black, Mean Annual Temperature, on Fahr. Scale.














A.K. JOHNSTON F.R.S.E.













OF THE MOST IMPORTANT IS, GRAINS, & FRUITS.

ONES OF CLIMATE, & MOISTURE.
A. K. JOHNSTON, F.R.S.E.




1











The only **ATLAS** for which a **MEDAL** was awarded at
the International Exhibition of 1882.

Complete in One Volume, Imperial Folio, half-bound in russia or morocco, price £5, 15s. 6d.

THE ROYAL ATLAS

OF MODERN GEOGRAPHY.

IN A SERIES OF ENTIRELY ORIGINAL AND AUTHENTIC MAPS.

By **ALEX. KEITH JOHNSTON, LL.D., F.R.S.E., F.R.G.S.,**

GEOGRAPHER TO THE QUEEN FOR SCOTLAND;
AUTHOR OF THE 'PHYSICAL ATLAS,' THE 'DICTIONARY OF GEOGRAPHY,' ETC.

WITH A COMPLETE INDEX TO EACH MAP, CONTAINING REFERENCES TO NEARLY 150,000 PLACES IN THIS ATLAS.

A NEW EDITION, BROUGHT UP TO THE PRESENT TIME, CONTAINING A NEW MAP
OF NORTH POLAR REGIONS.

CONTENTS.

Plate		With Index to 1530 Places.	Plate		With Index to 1832 Places.
1.	THE WORLD (IN HEMISPHERES), . . .		25.	BASIN OF THE BALTIC SEA, . . .	
2.	CHART OF THE WORLD ON MERCA- TOR'S PROJECTION, . . .	1840 "	26.	EUROPEAN RUSSIA, . . .	3099 "
3.	EUROPE, . . .	2120 "	27.	SOUTH-WEST RUSSIA, . . .	3757 "
4.	BASIN OF THE NORTH ATLANTIC OCEAN, . . .	1012 "	28.	ASIA, . . .	4101 "
5, 6.	ENGLAND (Two Sheets), . . .	10,723 "	29.	TURKEY IN ASIA (ASIA MINOR) and TRANSCASIA, . . .	2857 "
7, 8.	SCOTLAND (Two Sheets), . . .	9302 "	30.	PALESTINE, . . .	3104 "
9.	IRELAND, . . .	5328 "	31.	PERSIA and AFGHANISTAN, . . .	2153 "
10.	FRANCE, IN DEPARTMENTS, . . .	4565 "	32, 33.	INDIA (Two Sheets), . . .	7804 "
11.	SPAIN and PORTUGAL, . . .	4324 "	34.	CHINA and JAPAN, . . .	2510 "
12.	BASIN OF MEDITERRANEAN SEA, . . .	2193 "	35.	OCEANIA, . . .	2735 "
13, 14.	NORTHERN ITALY and SOUTHERN ITALY (Two Sheets), . . .	6349 "	36.	SOUTH AUSTRALIA, NEW SOUTH WALES, and VICTORIA, . . .	2769 "
15.	SWITZERLAND, and the ALPS of SAVOY and PIEDMONT, . . .	5014 "	37.	AFRICA, . . .	4048 "
16.	BELGIUM, . . .	2167 "	38.	NORTH-WESTERN AFRICA, } One SOUTHERN AFRICA, } Sheet, {	1519 "
17.	THE NETHERLANDS, } One Sheet, { DENMARK and ICELAND, } 2284 "	2284 "	39, 40.	EGYPT, NUBIA, ABYSSINIA, and ARABIA PETRÆA (Two Sheets), . . .	1261 "
18, 19.	HANOVER, BRUNSWICK, } One MECKLENBURG, and OLD- } Sheet, {	2037 "	41.	NORTH AMERICA, . . .	3166 "
20, 21.	EMPIRE OF GERMANY, . . .	1193 "	42, 43.	CANADA, NEW BRUNSWICK, NOVA SCOTIA, and NEWFOUNDLAND (Two Sheets), . . .	2791 "
22.	AUSTRIAN EMPIRE (Two Sheets), . . .	7176 "	44, 45.	UNITED STATES OF NORTH AMERICA (Two Sheets), . . .	3203 "
23.	TURKEY IN EUROPE, . . .	6412 "	46.	WEST INDIES and CENTRAL AMERICA, . . .	6094 "
24.	GREECE and the IONIAN ISLANDS, . . .	2209 "	47, 48.	SOUTH AMERICA, . . .	1188 "
	SWEDEN and NORWAY (SCANDINAVIA), 1691 "	2188 "		NORTH POLAR CHART, . . .	5479 "

Each Plate may be had separately, with its Index, price 3s.

OPINIONS OF THE PRESS.

Athenæum.

Under the name of *The Royal Atlas of Modern Geography*, Messrs Blackwood and Sons have published a book of maps, which, for care of drawing and beauty of execution, appears to leave nothing more to hope for or desire. Science and art have done their best upon this magnificent book. Mr A. Keith Johnston answers for the engraving and printing: to those who love clear forms and delicate bold type we need say no more. All that maps should be, these maps are; honest, accurate, intelligible guides to narrative or description. A very good feature is the Index of Names attached to each plate, with the easy mechanical arrangement of lines by which any particular name that may be sought is at once found on the map. Of the many noble atlases prepared by Mr Johnston and published by Messrs Blackwood and Sons, this Royal Atlas will be the most useful to the public, and will deserve to be the most popular.

Examiner.

There has not, we believe, been produced for general public use a book of maps equal in beauty and completeness to the Royal Atlas just issued by Mr A. K. Johnston. . . . In beauty and clearness of engraving it has never been surpassed; and the fulness of information in such maps as the five representing the United Kingdom, or the two representing India, is marvellous.

KEITH JOHNSTON'S ROYAL ATLAS.

OPINIONS OF THE PRESS—*Continued.*

The Times.

No one can look through Mr Keith Johnston's Atlas without seeing that it is the best which has ever been published in this country. To those who have a mean idea of our British map-makers perhaps this is not saying much. Let us therefore add that the new Atlas takes away from us a reproach, and is worthy of a place beside the best of the Continental ones. It is a fit successor to that other production of Mr Johnston's, the *Physical Atlas*, which is quite unique among such works for beauty of execution and richness of information. In the present volume, we have, at a charge of five and a half guineas, about fifty most carefully prepared and highly finished maps, together with a special index to each, containing all the names that appear in it. The price, therefore, of these remarkably accurate and beautiful maps is no more than 2s. for each—a fabulous sum, if we consider the amount of work which they contain.

Saturday Review.

The completion of Mr Keith Johnston's *Royal Atlas of Modern Geography* claims a special notice at our hands. While Mr Johnston's maps are certainly unsurpassed by any for legibility and uniformity of drawing, as well as for accuracy and judicious selection, this eminent geographer's Atlas has a distinguishing merit in the fact that each map is accompanied by a special index of remarkable fulness. The labour and trouble of reference are in this way reduced to a minimum. . . . The number of places enumerated in the separate indices is enormous. We believe, indeed, that every name which appears in the maps is registered in the tables; and as each place is indicated by two letters, which refer to the squares formed by the parallels of latitude and longitude, the method of using the index is extremely easy and convenient. . . . We know no series of maps which we can more warmly recommend. The accuracy, wherever we have attempted to put it to the test, is really astonishing.

Guardian.

This is, beyond question, the most splendid and luxurious, as well as the most useful and complete, of all existing atlases. To a habitual consulter of maps (which every reader of history and every man of science ought to be) there can hardly be a greater luxury than these beautiful sheets, which it is a pleasure for the eye to travel over, and upon which a vast quantity of the finest and most delicate work represents a more than corresponding mass of various information. In one large but not cumbrous folio volume Mr Keith Johnston has given us as perfect an atlas as can well be desired, embracing the results of all the most recent discoveries in every part of the world, as well as most of the territorial changes that have lately occurred in Europe. . . . A close examination of several different parts of Europe which are very familiar to us, satisfies us of the remarkable accuracy with which the work is done; it is very rarely that we miss even a new road, where, according to the scale of the map, a road should have been given.

Scotsman.

To go over the maps *seriatim*, noticing additions and improvements, would be tedious and somewhat unprofitable. An almost daily reference to, and comparison of it with others, since the publication of the First Part some two years ago until now, enables us to say, without the slightest hesitation, that this is by far the most complete and authentic atlas that has yet been issued. In the whole of that time we failed to detect more than two or three errors, and these were so very trifling that we have forgotten what they were.

In One Volume, Imperial Octavo, pp. 680, price 21s.

INDEX GEOGRAPHICUS:

BEING A LIST, ALPHABETICALLY ARRANGED, OF THE PRINCIPAL PLACES ON THE GLOBE, WITH THE COUNTRIES AND SUBDIVISIONS OF THE COUNTRIES IN WHICH THEY ARE SITUATED, AND THEIR LATITUDES AND LONGITUDES.

COMPILED SPECIALLY WITH REFERENCE TO KEITH JOHNSTON'S ROYAL ATLAS, BUT APPLICABLE TO ALL MODERN ATLASES AND MAPS.

In the preparation of this work, two objects have been specially kept in view: the first, to render it as completely as possible an Index of easy reference to the 'Royal Atlas'; and the second, to make it available not only for other Atlases, but to a great degree as a substitute for a Gazetteer.

In carrying out the first of these objects, care has been taken to secure insertion of every name that occurs in the 'Royal Atlas', which contains a larger number of positions than can be found in any similar work hitherto published. The original references to the Maps of that Atlas by means of Letters designating spaces on them are still retained; but, in addition to this, the Latitude and Longitude of every place is now given, by means of which the Index can be applied to any other geographical work.

With a view to the promotion of the general usefulness of the Index as a work of general reference, not only have all the names as they appear in the 'Royal Atlas' been inserted in it, but, so far as they could be collected, all popular variations of these as regards spelling, all historical forms or names, and in general every form under which they are likely to occur in other works, have also been included. Thus, while "Puducheri" and "Kachh" are adopted in the Map as more in consonance with Eastern pronunciation than "Pondichery" and "Cutch," the latter, as of constant recurrence in ordinary usage, are likewise given in the Index.

In addition to the Countries or Kingdoms, the subdivision of these countries in which the place is situated—as Counties, Provinces, Districts, Presidencies, &c.—has been appended to every name; and it is hoped that this will materially enhance its value for all ordinary purposes of general reference.

'By far the most complete list of the names of places in existence.'—*Address of the President of Royal Geographical Society.*

"It is, in fact, a Gazetteer of unparalleled fulness."—*Spectator.*

EDUCATIONAL WORKS.

ZOOLOGY.

- OUTLINES OF NATURAL HISTORY**, for Beginners; being Descriptions of a Progressive Series of Zoological Types. Fifty-two Engravings. 1s. 6d.
- INTRODUCTORY TEXT-BOOK OF ZOOLOGY**, for the Use of Junior Classes. By the same Author. With 127 Engravings. New Edition. 2s. 6d.
- TEXT-BOOK OF ZOOLOGY**, for the Use of Schools. By the same Author. Second Edition, enlarged. Crown 8vo, with 188 Engravings on Wood. 6s.
- A MANUAL OF ZOOLOGY**, for the Use of Students. With a General Introduction on the Principles of Zoology. By HENRY ALLEYNE NICHOLSON, M.D., F.R.S.E., F.G.S., &c., Professor of Natural History in the University of Toronto. Third Edition. Crown 8vo, pp. 706, with 280 Engravings on Wood, 12s. 6d.
- INTRODUCTION TO THE STUDY OF BIOLOGY**. By the same Author. Crown 8vo, with numerous Illustrations. 5s.

METEOROLOGY.

- INTRODUCTORY TEXT-BOOK OF METEOROLOGY**. By Alexander Buchan, M.A., F.R.S.E., Secretary of the Scottish Meteorological Society, Author of 'Handy Book of Meteorology,' &c. Crown 8vo, with 8 Coloured Charts and other Engravings, pp. 218. 4s. 6d.

GEOLOGY.

- INTRODUCTORY TEXT-BOOK OF GEOLOGY**. By David Page, LL.D., Professor of Geology in the Durham University of Physical Science, Newcastle. With Engravings on Wood and Glossarial Index. Tenth Edition. 2s. 6d.
- ADVANCED TEXT-BOOK OF GEOLOGY**, Descriptive and Industrial. By the Same. With Engravings, and Glossary of Scientific Terms. Fifth Edition, revised and enlarged. 7s. 6d.
- THE GEOLOGICAL EXAMINATOR**. A Progressive Series of Questions adapted to the Introductory and Advanced Text-Books of Geology. Prepared to assist Teachers in framing their Examinations, and Students in testing their own Progress and Proficiency. By the Same. Fifth Edition. 9d.
- GEOLOGY FOR GENERAL READERS**. A Series of Popular Sketches in Geology and Palaeontology. By the Same. Third Edition, enlarged. 6s.
- HANDBOOK OF GEOLOGICAL TERMS, GEOLOGY, AND PHYSICAL GEOGRAPHY**. By the Same. Second Edition, enlarged. 7s. 6d.
- CHIPS AND CHAPTERS**. A Book for Amateurs and Young Geologists. By the Same. 5s.
- THE PAST AND PRESENT LIFE OF THE GLOBE**. With numerous Illustrations. By the Same. Crown 8vo. 6s.
- THE EARTH'S CRUST: A Handy Outline of Geology**. By the Same. Sixth Edition. 1s.
- SYNOPSIS OF SUBJECTS** taught in the Geological Class, College of Physical Science, University of Durham, Newcastle-on-Tyne. By the Same. Fcap., cloth, 2s. 6d.

PHYSICAL GEOGRAPHY.

- INTRODUCTORY TEXT-BOOK OF PHYSICAL GEOGRAPHY**. With Sketch-Maps and Illustrations. By DAVID PAGE, LL.D., Professor of Geology in the Durham University of Physical Science, Newcastle. Sixth Edition. 2s. 6d.
- ADVANCED TEXT-BOOK OF PHYSICAL GEOGRAPHY**. By the Same. With Engravings. Second Edition, enlarged. 5s.
- EXAMINATIONS ON PHYSICAL GEOGRAPHY**. A Progressive Series of Questions, adapted to the Introductory and Advanced Text-Books of Physical Geography. By the Same. 9d.
- "Few of our handbooks of popular science can be said to have greater or more decisive merit than those of Mr Page on Geology and Palaeontology. They are clear and vigorous in style, they never oppress the reader with a pedantic display of learning, nor overwhelm him with a pompous and superfluous terminology; and they have the happy art of taking him straightway to the face of nature herself, instead of leading him by the tortuous and bewildering paths of technical system and artificial classification."*—Saturday Review.

- COMPARATIVE GEOGRAPHY**. By Carl Ritter. Translated by W. L. Gage. Fcap., 3s. 6d.

BOTANY.

- ADVANCED TEXT-BOOK OF BOTANY**. For the Use of Students. By Robert BROWN, M.A., Ph.D., F.R.G.S., Lecturer on Botany under the Science and Art Department of the Committee of the Privy Council on Education. [In the Press.]

PHYSICS.

- ELEMENTARY HANDBOOK OF PHYSICS**. With 210 Diagrams. By William ROSSITER, F.R.A.S., &c. Crown 8vo, pp. 390. 5s.

WILLIAM BLACKWOOD & SONS, EDINBURGH AND LONDON.

EDUCATIONAL WORKS—*Continued.*

ARITHMETIC, &c.

- THE THEORY OF ARITHMETIC.** By David Munn, F.R.S.E., Mathematical Master Royal High School of Edinburgh. Crown 8vo, pp. 294. 5s.
- ELEMENTARY ARITHMETIC.** By Edward Sang, F.R.S.E. This Treatise is intended to supply the great desideratum of an intellectual instead of a routine course of instruction in Arithmetic. Post 8vo. 5s.
- THE HIGHER ARITHMETIC.** By the same Author. Being a Sequel to 'Elementary Arithmetic.' Crown 8vo. 5s.
- FIVE-PLACE LOGARITHMS.** Arranged by E. Sang, F.R.S.E. Sixpence, for the Waistcoat-Pocket.
- TREATISE ON ARITHMETIC, with numerous Exercises for teaching in Classes.** By JAMES WATSON, one of the Masters of Heriot's Hospital. Foolscep. 1s.
- A GLOSSARY OF NAVIGATION.** Containing the Definitions and Propositions of the Science, Explanation of Terms, and Description of Instruments. By the Rev. J. B. HARBORD, M.A., Assistant Director of Education, Admiralty. Crown 8vo. Illustrated with Diagrams. 6s.
- DEFINITIONS AND DIAGRAMS IN ASTRONOMY AND NAVIGATION.** By the Same. 1s. 6d.
- EXAMPLES IN BOOK-KEEPING.** A Treatise specially adapted for Schools. By the Rev. JOHN CONSTABLE, M.A., Principal of the Royal Agricultural College, Cirencester. [*In the Press.*]

ENGLISH LANGUAGE.

- ETYMOLOGICAL AND PRONOUNCING DICTIONARY OF THE ENGLISH LANGUAGE.** Including a very Copious Selection of Scientific Terms. For use in Schools and Colleges, and as a Book of General Reference. By the Rev. JAMES STORMONTH. The Pronunciation carefully revised by the Rev. P. H. PHELP, M.A. Cantab. Crown 8vo, pp. 758. 7s. 6d.
- THE SCHOOL ETYMOLOGICAL DICTIONARY AND WORD-BOOK:** combining the Advantages of an Ordinary School Dictionary and an Etymological Spelling-Book. By the same Author. Foolscep 8vo. 2s.
- ENGLISH PROSE COMPOSITION: A Practical Manual for Use in Schools.** By James CURRIE, M.A., Principal of the Church of Scotland Training College, Edinburgh. Ninth Edition. 1s. 6d.
- PROGRESSIVE AND CLASSIFIED SPELLING-BOOK.** By Hannah R. Lockwood, Authoress of 'Little Mary's Mythology.' Fcap. 8vo. 1s. 6d.
- A MANUAL OF ENGLISH PROSE LITERATURE, Biographical and Critical; designed mainly to show characteristics of style.** By W. MINTO, M.A. Crown 8vo. 10s. 6d.
- A TREASURY OF THE ENGLISH AND GERMAN LANGUAGES.** Compiled from the best Authors and Lexicographers in both Languages. Adapted to the Use of Schools, Students, Travellers, and Men of Business; and forming a Companion to all German-English Dictionaries. By JOSEPH CAUVIN, LL.D. & PH.D., of the University of Göttingen, &c. Crown 8vo. 7s. 6d. bound in cloth.
- CHOIX DES MEILLEURES SCENES DE MOLIERE, avec des Notes de Divers Commentateurs, et autres Notes Explicatives.** Par DR^E E. DUBUC. Fcap. 8vo. 4s. 6d.

HISTORY.

- EPITOME OF ALISON'S HISTORY OF EUROPE, for the Use of Schools.** Sixteenth Edition. Post 8vo, pp. 604. 7s. 6d. bound in leather.
- ATLAS TO EPITOME OF THE HISTORY OF EUROPE.** Eleven Coloured Maps. By A. KEITH JOHNSTON, LL.D., F.R.S.E. In 4to. 7s.
- THE EIGHTEEN CHRISTIAN CENTURIES.** By the Rev. James White, Author of 'The History of France.' School Edition, post 8vo, with Index. 6s.
- HISTORY OF FRANCE, from the Earliest Times.** By the Rev. James White, Author of 'The Eighteen Christian Centuries.' Post 8vo. 6s., with Index.
- FACTS AND DATES; or, The Leading Events in Sacred and Profane History, and the Principal Facts in the various Physical Sciences: the Memory being aided throughout by a Simple and Natural Method. For Schools and Private Reference.** By the Rev. ALEX. MACKAY, LL.D., F.R.G.S., Author of 'A Manual of Modern Geography,' &c. Second Edition, crown 8vo, pp. 336. 4s.

AGRICULTURE.

- CATECHISM OF PRACTICAL AGRICULTURE.** By Henry Stephens, F.R.S.E., Author of the 'Book of the Farm.' With Engravings. 1s
- PROFESSOR JOHNSTON'S CATECHISM OF AGRICULTURAL CHEMISTRY.** A New Edition, edited by PROFESSOR VOELCKER. With Engravings. 1s.
- PROFESSOR JOHNSTON'S ELEMENTS OF AGRICULTURAL CHEMISTRY AND GEOLOGY.** A New Edition, revised and brought down to the present time, by G. T. ATKINSON, B.A., F.C.S., Clifton College. Foolscep 8vo. 6s.

WILLIAM BLACKWOOD & SONS, EDINBURGH AND LONDON.

NOW PUBLISHING.

ANCIENT CLASSICS FOR ENGLISH READERS

BY VARIOUS AUTHORS.

Edited by Rev. W. LUCAS COLLINS, M.A.

Author of 'Etoniana,' 'The Public Schools,' &c.

OPINIONS OF THE PRESS.

"We gladly avail ourselves of this opportunity to recommend the other volumes of this useful series, most of which are executed with discrimination and ability."—*Quarterly Review*.

"It is difficult to estimate too highly the value of such a series as this in giving 'English readers' an insight, exact as far as it goes, into those olden times which are so remote and yet to many of us so close. It is in no wise to be looked upon as a rival to the translations which have at no time been brought forth in greater abundance or in greater excellence than in our own day. On the contrary, we should hope that these little volumes would be in many cases but a kind of stepping-stone to the larger works, and would lead many who otherwise would have remained in ignorance of them to turn to the versions of Conington, Worsley, Derby, or Lytton. In any case a reader would come with far greater knowledge, and therefore with far greater enjoyment, to the complete translation, who had first had the ground broken for him by one of these volumes."—*Saturday Review*.

The Volumes published contain :—

I.—HOMER: THE ILIAD. By the Editor.

"We can confidently recommend this first volume of 'Ancient Classics for English Readers' to all who have forgotten their Greek and desire to refresh their knowledge of 'Homer.' As to those to whom the series is chiefly addressed, who have never learnt Greek at all, this little book gives them an opportunity which they had not before—an opportunity not only of remedying a want they must often have felt, but of remedying it by no patient and irksome toil, but by a few hours of pleasant reading."—*Times*.

II.—HOMER: THE ODYSSEY. By the Editor.

III.—HERODOTUS. By G. C. Swayne, M.A.

"This volume altogether confirms the highest anticipations that were formed as to the workmanship and the value of the series."—*Daily Telegraph*.

IV.—THE COMMENTARIES OF CÆSAR.

By ANTHONY TROLLOPE.

"We can only say that all admirers of Mr Trollope will find his 'Cæsar' almost, if not quite, as attractive as his most popular novel, while they will also find that the exigencies of faithful translation have not been able to subdue the charm of his peculiar style."—*Vanity Fair*.

V.—VIRGIL. By the Editor.

"Such a volume cannot fail to enhance the reputation of this promising series, and deserves the perusal of the most devoted Latinists, not less than of the English readers for whom it is designed."—*Contemporary Review*.

VI.—HORACE. By Theodore Martin.

"Though we have neither quoted it, nor made use of it, we have no hesitation in saying that the reader, who is wholly, or most part, unable to appreciate Horace untranslated, may, with the insight he gains from the lively, bright, and, for its size, exhaustive little volume to which we refer, account himself hereafter familiar with the many-sided charms of the Venesian, and able to enjoy allusions to his life and works which would otherwise have been a sealed book to him."—*Quarterly Review*.

VII.—ÆSCHYLUS. By R. S. Copleston, B.A.

"A really delightful little volume."—*The Examiner*.

VIII.—XENOPHON. By Sir Alex. Grant, Bart.

Principal of the University of Edinburgh.

"Sir Alexander Grant tells the story of Xenophon's life with much eloquence and power. It has evidently been with him a labour of love, while his wide reading and accurate scholarship are manifest on nearly every page."—*The Examiner*.

IX.—CICERO. By the Editor.

"No charm of style, no facility and eloquence of illustration, is wanting to enable us to see the great Roman advocate, statesman, and orator, in the days of Rome's grandeur, in the time of her first fatal hastening to her decadence, with whom fell her liberty two thousand years ago. The first lines of introduction to this fascinating book are full of help and light to the student of the classical times who has not mastered the classical literature, and in whose interests this book is done, simply to perfection."—*Saunders' News Letter*.

X.—SOPHOCLES. By C. W. Collins, M.A.

"Mr Collins shows great skill and judgment in analysing and discriminating the plays of the sweet singer of Colonus."—*Guardian*.

XI.—PLINY'S LETTERS. By Rev. A. Church M.A., and Rev. W. J. Brodribb, M.A.

"The writers of the volume before us have contrived, out of their intimate and complete familiarity with their subject, to place the man, his traits of character, his friends, and his surroundings, so vividly before us, that a hitherto shadowy acquaintance becomes a distinct and real personage."—*Saturday Review*.

XII.—EURIPIDES. By W. B. Donne.

"We take leave of a tribute to the merit of Euripides, which is calculated to enhance the appreciation of the most pathetic and philosophic of Greek dramatists."—*Saturday Review*.

XIII.—JUVENAL. By Edward Walford, M.A.

"This is one of the best executed volumes of the whole series of 'Ancient Classics,' and exhibits Mr Walford's critical powers in a very favourable light."—*Pall Mall Gazette*.

XIV.—ARISTOPHANES. By the Editor.

"No single volume has been better done than 'Aristophanes' by the Rev. W. Lucas Collins, who edits the whole of the series."—*Standard*.

XV.—HESIOD AND THEOGNIS. By the Rev. James Davies, M.A.

"If all the other works of this series were written with the same ability as Hesiod and Theognis, a very valuable addition, in an unpretending form, will have been made to the literature of the country."—*Saturday Review*.

XVI.—PLAUTUS AND TERENCE. By the Editor.

"One of the very best of this admirable series that has yet come under our notice. For the first time, indeed, the true meaning and purpose of the works of the two oldest Latin dramatists are thoroughly investigated, whilst the elegance of their phraseology and the unity of their design are equally exemplified."—*Bell's Weekly Messenger*.

XVII.—TACITUS. By W. B. Donne.

"Mr Donne's sketch of the life and writings of Tacitus is worthy of the excellent series of 'Ancient Classics' to which it belongs. It is evident that Mr Donne has studied his author deeply and thoroughly, at once recognising his defects and appreciating his merits. The little work before us is a brief but perspicuous summary of the works of the great historian, calculated not only to interest the English readers for whom it is primarily intended, but also to assist the student who is anxious to make himself acquainted with the history of the first century after Christ. We anticipate for the work a great success."—*Athenæum*.

XVIII.—LUCIAN. By the Editor.

Other Authors, by various contributors, are in preparation.

A Volume is published Quarterly, price 2s. 6d.

WILLIAM BLACKWOOD & SONS, EDINBURGH AND LONDON.

GEOGRAPHICAL AND EDUCATIONAL WORKS

PUBLISHED BY

W. & A. K. JOHNSTON,

GEOGRAPHERS AND ENGRAVERS TO THE QUEEN,

EDINBURGH AND LONDON;

Agents by Appointment for the Sale of the Ordnance Maps, the Geological Survey of Scotland, and Admiralty Charts.

THE NATIONAL SCHOOL BOARD SERIES OF LARGE WALL MAPS AND ILLUSTRATIONS.

By **KEITH JOHNSTON, LL.D.,**

GEOGRAPHER TO THE QUEEN FOR SCOTLAND.

This well-known Series possesses the following advantages:—

From the great demand, they are constantly at press, and no Map is ever printed without being thoroughly revised.

The Maps are printed by Steam, in permanent Oil Colours.

The Series is the most extensive published, consisting of upwards of 60 Maps and Illustrations, to which additions are constantly being made.

They are Mounted on Cloth and Rollers, Plain or Varnished, the best materials only being used.

The whole Series is of one uniform size—namely, 50 by 42 inches.

It is the cheapest ever published, the price being, Rollers Varnished, 12s.; Plain, 10s.

It is the *only Series* accompanied by Handbooks written expressly for each Map or Illustration. These are given *gratis* to purchasers.

It will be found **INDISPENSABLE** to **TEACHERS** and **MANAGERS** who require the **GOVERNMENT GRANT**.

Specimens and List sent gratis and post free to School Boards, Managers, and Teachers, on application to the Publishers.

GENERAL GEOGRAPHY.

EASTERN HEMISPHERE.

ENGLAND.
SCOTLAND.
IRELAND.
EUROPE.
ASIA.
AFRICA.
AMERICA.

CANAAN AND PALESTINE.
BRITISH ISLES.
FRANCE.
SPAIN.
ITALY.
CENTRAL EUROPE.
INDIA.

WESTERN HEMISPHERE.

NORTH AMERICA.
SOUTH AMERICA.
UNITED STATES.
CANADA, NOVA
SCOTIA, ETC.
AUSTRALIA.

NEW ZEALAND.
PACIFIC OCEAN.
TRAVELS OF ST PAUL.
WORLD, MERCATOR'S
PROJECTION.

EASTERN AND WESTERN HEMISPHERES (ONE MAP).

OUTLINE MAPS.

Uniform with the Maps of General Geography, showing the Outlines, Rivers, Mountains, Towns, and Political Divisions, *without names*.

WORLD, MERCATOR.
WORLD, IN HEMISPHERES.
EUROPE.

ASIA.
AFRICA.
AMERICA.

BRITISH ISLES.
ENGLAND.
SCOTLAND.

IRELAND.
INDIA.

CLASSICAL GEOGRAPHY.

ORBIS VETERIBUS
NOTIS.

ITALIA ANTIQUA.

GRÆCIA ANTIQUA.

ASIA MINOR.
ORBIS ROMANUS.

PHYSICAL GEOGRAPHY.

WORLD, IN HEMISPHERES. | EUROPE. | ASIA. | AFRICA. | AMERICA.

Also now ready, **PHYSICAL CHART OF THE WORLD**, on MERCATOR'S PROJECTION, suitable for School or Library.
50 by 42 inches. Price £1. 1s.

Each Physical Map is accompanied by a Handbook, with a Coloured Sketch-Map.

SET OF MAPS IN STAND.

THE STAND is mounted on Castors, and contains 10 Coloured Maps on a continuous web of cloth, which revolves vertically over Rollers by turning the handle at the side, so that the Maps are exhibited in rotation.

LIST OF THE MAPS.

E. Hemisphere—W. Hemisphere—Europe—Asia—Africa—America—England—Scotland—Ireland, & Palestine.

The Maps measure 4 feet 2 inches by 3 feet 6 inches. The Stand is 6 feet 6 inches high by 4 feet 7 inches broad. It can be taken in pieces, packed in matting, and sent with safety to any part of the country.

Price of the 10 Maps and Painted Stand, £7. 17s. 6d.

Black Board behind for Arithmetic or Diagrams, 10s. 6d. extra.

SET OF MAPS IN CASE.

THE CASE for hanging on a wall contains 10 Coloured Maps on Cloth and Rollers, and is so constructed that any Map can be drawn down as required, and pulled up again by the Cords at the side.

LIST OF THE MAPS.


E. Hemisphere—W. Hemisphere—Europe—Asia—Africa—America—England—Scotland—Ireland, & Palestine.

The Maps measure 4 feet 2 inches by 3 feet 6 inches. The Case is 4 feet 8 inches long by 1 foot 9 inches high; and as it is only 4 inches thick, it projects very little from the wall. It can be packed in matting, and sent with safety to any part of the country.

Price of the 10 Maps and Painted Case, £7. 7s.

ILLUSTRATIONS OF NATURAL PHILOSOPHY.

- No. I.
PROPERTIES OF BODIES,
37 Diagrams.
- No. II.
MECHANICAL POWERS,
47 Diagrams.
- No. III.
HYDROSTATICS,
28 Diagrams.
- No. IV.
HYDRAULICS,
27 Diagrams.
- No. V.
HUMAN ANATOMY,
No. 1, 27 Diagrams.
- No. VI.
HUMAN ANATOMY,
No. 2, 42 Diagrams.
- No. VII.
STEAM-ENGINES AND BOILERS,
15 Diagrams.



- No. VIII.
BOTANY,
Sheet 1.
36 Illustrations.
- No. IX.
BOTANY,
Sheet 2.
36 Illustrations.
- No. X.
BOTANY,
Sheet 3.
36 Illustrations.
- No. XI.
BOTANY,
Sheet 4.
44 Illustrations.
- In preparation,
ASTRONOMY.

Uniform in Size and Price with the National School Board Series of Large Wall Maps.

SIZE—4 FEET 2 INCHES BY 3 FEET 6 INCHES.

PRICE—Coloured, on Cloth and Roller, 10s.
" " " Varnished, 12s.

A Handbook is given free with each Wall Map and Illustration.

METRIC SYSTEM.

A SYNOPSIS TABLE, WITH DIAGRAMS, SHOWING THE ACTUAL SIZE OF WEIGHTS AND MEASURES,
BY C. H. DOWLING, C.E., WITH HANDBOOK BY JAMES YATES, Esq., M.A., & C. & C.
Size, 5 feet by 4 feet 2 inches. Price, coloured, on Cloth and Rollers, 15s.; varnished, 17s.; in 4to Portfolio, 15s.

Also now ready,

A CHEAP EDITION of the Above,

In which the Weights and Measures are shown one-fourth of the Actual Size.
Size, 33 by 27 inches. Price, coloured, on Cloth and Rollers, 5s., varnished, 6s.

SMALL WALL MAPS.

EASTERN HEMISPHERE.	ENGLAND.	AFRICA.
WESTERN HEMISPHERE.	SCOTLAND.	AMERICA.
E. & W. HEMISPHERES (1 Map).	IRELAND.	CANAAN AND PALESTINE.
WORLD (Mercator).	EUROPE.	CANADA, UNITED STATES, AND
BRITISH ISLES.	ASIA.	MEXICO.

Size, 33 by 27 inches. Price, coloured, on Cloth and Rollers, 5s.; varnished, 6s.

Uniform in Size, Price, and Styles of Mounting, with the Small Wall Maps.

A MAP ILLUSTRATIVE OF GEOGRAPHICAL TERMS, WITH GLOSSARY.

CHRONOLOGICAL CHART OF ANCIENT HISTORY, from the Creation to the Fall of the Western Roman Empire, with GLOSSARY.

10 of the above MAPS, mounted on a Stand (see preceding page), with Diagram Board on back, £5, 5s. The Stand measures 6 feet in height by 3 feet 3 inches in breadth.

In Case (see preceding page), the 10 MAPS cost £4, 14s. 6d. Size of the Case, 3 feet 3 inches by 1 foot 6 inches.

JOHNSTON'S MODERN GLOBES, WITH ALL THE MOST RECENT DISCOVERIES.

30-INCH TERRESTRIAL GLOBE, Black Stand, £8 18 6	12-INCH GLOBE, Black Stand, £2 12 6
18-INCH GLOBE, Low Black Stand, 5 5 0	" " High Mahogany Stand with
" " Mahogany Stand, 5 15 6	Compass, 3 13 6
" " High " " 7 17 6	6-INCH GLOBE, Low Black Stand, 1 5 0
	3-INCH GLOBE, Rosewood Stand, 0 8 6

When packing-cases are returned and received, a small charge only is made for the use of them. The Publishers do not, under any circumstances, hold themselves responsible for any damage Globes may sustain after leaving their premises.

LARGE OUTLINE CHARTS OF THE WORLD.

No. I., in 20 sheets—size, when joined, 15 feet by 12 feet; price in Sheets, plain, £5—or mounted in 3 pieces to join up, £7. No. II., in 4 sheets—size, when joined, 6 feet by 4 feet 4 inches; price in Sheets, plain, 18s.—or on Cloth and Roller, 30s.

JOHNSTON'S CHART OF THE WORLD ON MERCATOR'S PROJECTION.

Size, 6 feet by 4 feet 8 inches. Price on Roller varnished, or in quarto morocco Case, £3, 3s.

LARGE SCHOOL-MAP OF EUROPE, on Roller, Varnished.

Size, 5 feet 8 inches by 5 feet 3 inches.

men-Sheet post free on application.

